item 10 Number	03837 Net Scanned
Author	De Sylva, Donald P.
Corporate Author	University of Miami, Rosenstiel School of Marine and At
Report/Article Title	The Effects of Herbicides in South Vietnam: Part B, Working Papers, February 1974: Effects of Mangrove Defoliation on the Estuarine Ecology and Fisheries of South Vietnam
Journal/Book Title	
Year	1974
Month/Bay	February
Color	
Number of Images	128
Descripton Notes	AD-779 014. Contribution No. 1702. Last three pages are duplicated.

.

JML \$4.75

AD-779 014

THE EFFECTS OF HERBICIDES IN SOUTH VIETNAM PART B. WORKING PAPERS: EFFECTS OF MANGROVE DEFOLIATION ON THE ESTUARINE ECOLOGY AND FISHERIES OF SOUTH VIETNAM

NATIONAL ACADEMY OF SCIENCES-NATIONAL RESEARCH COUNCIL

FEBRUARY 1974

DISTRIBUTED BY:

National Technical Information Service U. S. DEPARTMENT OF COMMERCE

AD-779 014

THE EFFECTS OF HERBICIDES IN SOUTH VIETNAM

PART B: WORKING PAPERS

FEBRUARY 1974

Effects of Mangrove Defoliation on the Estuarine Ecology and Fisheries of South Vietnam

DONALD P. DE SYLVA AND HARDING B. MICHEL

Reproduced by NATIONAL TECHNICAL INFORMATION SERVICE U S Department of Commerce Springfield VA 22151

NATIONAL ACADEMY OF SCIENCES - NATIONAL RESEARCH COUNCIL

WASHINGTON, D.C. 20418

DISTRIBUTION STATEMENT A

Approved for public release; Distribution Unlimited

Effects of Mangrove Defoliation on the Estuarine Ecology and Fisheries of South Vietnam^a

DONALD P. DE SYLVA AND HARDING B. MICHEL^b

Mangroves are an intrinsic part of the tropical coastal and estuarine environment (Lauff 1967). Although some species are restricted to land, a large part of the mangrove ecosystem comprises aquatic habitats characterized by the ebb and flow of tidal salt water (Macnae 1968). The complex aerial root system acts as substrate for the attachment of various invertebrate organisms such as worms, snails, oysters, mussels, and barnacles (Berry 1964). Beneath the water and submerged among the labyrinth of roots, the young stages of fishes, crabs, and shrimps find food and shelter from predators. Thus, the mangroves act as nursery grounds for a variety of aquatic organisms (Fischer-Piette 1931).

The root complex of mangroves also serves as a sediment trap to keep soil from eroding (Stephens 1963), and silt-laden water flowing across mangroves is filtered by the roots (Gledhill 1963, Gigliogli and Thornton 1965). The result is that the water becomes clearer (Tabb et al. 1962), permitting greater light penetration, and therefore photosynthesis, to occur at levels deeper in the water column than would be possible in turbid water. Hence, production of aquatic organisms is indirectly increased by mangroves (Davis 1940), which are among the most fertile ecosystems (Golley et al. 1962).

^aContribution No. 1702, University of Miami, Rosenstiel School of Marine and Atmospheric Science.

^bDrs. de Sylva and Michel, consultants to the Committee on the Effects of Herbicides in Vietnam, are affiliated with the Rosenstiel School of Marine and Atmospheric Science, University of Miami, 10 Rickenbacker Causeway, Miami, Florida 33149.

Recently it has been found that the great worth of mangroves to the aquatic ecosystem is through the contribution of decaying leaves to the food chain in the aquatic environment (Heald 1971, Odum 1970, Odum and Heald 1972). As the leaves decompose in the water, they are changed biochemically by fungi into protein for various kinds of benthos and plankton. This organic debris amounts to approximately 3 tons/acre of potential food for plankton and benthos. The organic contribution from decomposing mangrove seedlings may be even greater. The food thus produced by fungi from these leaves is eaten by shrimps, crabs, worms, roundworms, and snails, as well as by several hundred different kinds of organisms in the plankton (Fell and Master 1973). In turn, these consumers are a primary source of food for tropical commerical fishes such as flounders, herrings, mullets, porgies, jacks, anchovies, snappers, and croakers (Freise 1935). Thus, destruction of mangroves reduces both food and shelter for important fishes, as well as invertebrates such as crabs and shrimps (Freyberg 1930).

On the basis of the above concepts, scientists have suggested that the destruction of large parts of the estuarine and coastal mangrove forests of South Vietnam (SVN) through military defoliation (Tschirley 1969, Odum 1971) should affect the life cycles of fishes dependent upon this region (Orians and Pfeiffer 1970, Erlich and Erlich 1972). Further, they expect a decrease in the commerical fishery for aquatic species that depend on the mangroves for food, shelter, or spawning grounds (Gerlach 1958, Kiener 1966, Walsh 1967, de Sylva 1970). Herbicides used for defoliation may have affected the fishes and invertebrates of the mangrove region either by killing them outright or by interfering, sublethally,

with normal physiological processes such as feeding, reproduction, behavior, and migratory patterns. Finally, it has been theorized (Meselson et al. 1970) that the herbicides--including their breakdown products, or contaminants accidentally produced during their manufacture-may have been passed on through the food chain by way of decaying mangrove leaves, and subsequently into water and sediments, plankton, intermediate-size predators, and thence finally into large fishes and invertebrates of commerical value which are so important in the Vietnamese diet. One such contaminant--dioxin--has been found in fish and shrimp taken near defoliated areas (Shapley 1973).

It was the purpose of this study, then, to attempt to evaluate some of the above factors by collecting data in the defoliated mangrove forest of SVN (Figures 1 and 2) and to compare ecological information with data from a mangrove region that had not been sprayed with herbicides (control region). Positions for collecting stations for studies made in the defoliated and nondefoliated areas are given in Table I.

The original intent was to select a control area in the Ca-Mau Peninsula, which has been relatively untouched by military defoliants, and which is distant from large urban populations or other sources of pollution. Security restrictions prevented us from entering this region, and a less desirable control area (subsequently referred to as the Vung-Tau Region) was necessarily selected. However, the Vung-Tau Region, even though it was the least sprayed mangrove area available to us, nevertheless had been subjected to herbicidal spraying in two contiguous regions in Phuoc-Tuy Province (Table II; Figures 3 and 4, Targets 1 and 2)



FIG. 1. Location of studies on the estuarine ecology and fisheries of SVN.

>



FIG. 2. Location of collecting stations in the Rung-Sat Special Zone and the Vung-Tau Region, SVN, 1972 and 1973.

.

Location of collecting stations (VND) in the Rung-Sat Special Zone and Vung-Tau Region, SVN, October-November 1972 and January 1973.

.

Station	Latitude	Longitude		Station	Latitud.	Longitude	
VND 10-11 = VND 66-67	10°40'06"N	106 ⁰ 46'38"E	Rung Sat	VND 40-41	10 ⁰ 27 * 09" N	107 ⁰ 11'08"E	Vung Tau
VND 12-13 = VND 64-65	10 ⁰ 38*05"N	106 ⁰ 49'00"E	Rung Sat	VND 42-43	10 ⁰ 28124"N	107 ⁰ 10 '13''E	Vung Tau
VND 14-15 = VND 62-63	10 ⁰ 36'34"N	106 ⁰ 52'02"E	Rung Sat	VND 44-45	10 ⁰ 27 ' 55"N	107 ⁰ 10'15"E	Vung Tau
VND 16-17 = VND 60-61	10 ⁰ 34 '50"N	106 ⁰ 50'46"E	Rung Sat	VND 46-47	10 ⁰ 27'14"N	107 ⁰ 10 ' 19 ''E	Vung Tau
VND 18-19 = VND 80-81	10 ⁰ 32 ' 25"N	106 ⁰ 50 '59'' E	Rung Sat	VND 48-49	10 ⁰ 27'17"N	107 ⁰ 09*35"E	Vung Tau
VND 20-21 = VND 72-73	10 ⁰ 31'22''N	106°52'32"E	Rung Sat	VND 50-51 = VND 86-87	10 ⁰ 27'30"N	107 ⁰ 08'49"E	Vung Tau
VND 22-23 = VND 70-71	10°30'40"N	106°57'17"E	Rung Sat	VND 52-53 = VND 88-89	10 ⁰ 27 ' 18"N	107°07'58"E	Vung Tau
VND 24-25 = VND 68-69	10 ⁰ 31'50"N	106 ⁰ 55'11"E	Rung Sat	VND 54-55 = VND 90-91	10 ⁰ 26'12"N	107°07'00"E	Vung Tau
VND 26-27	10 ⁰ 26 ' 47 "N	106 ⁰ 59†13"E	Rung Sat	VND 56-57	10 ⁰ 26'00"N	107 ⁰ 09'47"E	Vung Tau
VND 28-29 = VND 76-77	10 ⁰ 28'10"N	106°56'35"E	Rung Sat	VND 58-59 = VND 104-105	10 ⁰ 24 ' 52"N	107°08'54"E	Vung Tau
VND 30-31	10 ⁰ 28 ' 00"N	106 ⁰ 52"20"E	Rung Sat	VND 82-83	10 ⁰ 26'17"N	107 ⁰ 09'15"E	Vung Tau
VND 32-33 = VND 74-75	10 ⁰ 33'30"N	106 ⁰ 54 ' 05"E	Rung Sat	VND 84-85	10 ⁰ 27'51"N	107°09'36"E	Vung Tau
VND 34-35	10 ⁰ 27 '53"N	106 ⁰ 58'26"E	Rung Sat	VND 92-93	10 ⁰ 24 '49"N	107°05'45"E	Vung Tau
VND 78-79	10 ⁰ 28 ' 20"N	106 ⁰ 54'40"E	Rung Sat	VND 94-95	10 ⁰ 22'50"N	107 ⁰ 02'41"E	Vung Tau
VND 36-37 = VND 98-99	10 ⁰ 23'54"N	107 ⁰ 04'10"E	Vung Tau	VND 96-97	10 ⁰ 23'15"N	107 ⁰ 03'28"E	Vung Tau
VND 38-39 = VND 102-103	10 ⁰ 24'31"N	107 ⁰ 06'53"E	Vung Tau	VND 100-101	10 ⁰ 23'34"N	107°05'20"E	Vung Tau

and the second sec

Table II.

Spray missions in Phuoc-Tuy Province (near Vung-Tau), 1962-1968. Data from the List of 202 Task Realized file and HERBS tape.

		AGENT IN		
YEAR	PURPLE	ORANGE	WHITE	TOTAL
Target 1				
(and vicinity)				
1962	4,000 ⁸		*** 88	4,000
1964		14,000		14,000
1965	~~	5,000	***	5,000
1966		21,650		21,650
1967	APP 46 +	16,200	6,100	22,300
1968		11,675	45,000	56,675
TOTAL	4,000 ^a	68,525	51,100	123,625
Target 2				
1965		6,000		6,000
1966		21,600		21,600
1968			7,000	7,000
TOTAL		27,600	7,000	34,600
Target 3				
1965	مجه جغ	. 8,850	-	8,850
1966		5,150		5,150
1967		23,740		23,740
1968		20,650	29,400	50,050
TOTAL		58,390	29,400	87,790

^aTest spraying along Route 15.





as recently as 1968. The defoliating agents used were Orange (2,4-D and 2,4,5-T), Purple (same as Orange, with slight molecular variation), and White (2,4-D and picloram), totalling about 158,000 gal sprayed adjacent to the Vung-Tau Region (Table II). In comparison, the heavily defoliated Rung-Sat Special Zone received direct application of over 1.1 million gal of spray during a comparable period (Table III). Target areas 1 and 2 comprise hilly areas, and during the rainy season it would be expected that any residual herbicides, contaminants, or breakdown products from these targets could be carried into the mangrove estuary of the Vung-Tau Region. Another factor is that the two areas are not strictly comparable in their soils, muds, and tidel characteristics, as will be discussed.

Limitations of Study

A number of factors clearly limit the conclusions that can be drawn based upon the present study. Chief among these is the lack of almost any qualitative or quantitative background data both in the defoliated area and the control region. Nearly all aquatic studies in Indochina seem to have been confined to open waters or inland streams and ponds (Soulier 1963, de Sylva 1973a); virtually nothing is known about the estuarine ecology of mangroves in SVN or the organisms inhabiting these regions. Some conclusions may be inferred based on the <u>kinds</u> of animals living there, by using studies following the defoliation, and comparing them with what is known about similar mangrove habitats in Malaysia, Africa, or the Caribbean, because usually the same <u>families</u> of animals inhabit the same kinds of mangrove communities. However, hypothetical data cannot be extrapolated on the numbers of aquatic animals

Table III.

Defoliation missions in the Rung-Sat Special Zone. Data from the List of 202 Task Realized file and HERBS tape.

	Orai	nge	. <u>Whi</u>	te	Blue				
Year	Missions	<u>Gallons</u>	Missions	<u>Gallons</u>	Missions	<u>Gallons</u>			
1 9 64		31,650 ⁸	· •						
1965		27,000 ^b		er	u	ite -st.			
1966	32	74,900	5	16,625	3	8,100			
1 9 67	89	315,200	17	46,235	4	13,500			
1968	39	161,850	66	245,325	10	27,500			
1969	22	101,300	4	26,200		-			
1970	5	12,400	3	9,000					
Total	187	724,300	95	343,385	17	49,100			

^aFrom List of 202 Task Realized File

^bFrom HERBS tape

ų,

living in Vietnam prior to mangrove defoliation. Thus, we really do not know what kinds or how many organisms were there before wartime conditions prevailed. Large-scale defoliation or other alteration of an environment would be expected to produce widespread changes in the kinds of animals and plants because the physical environment is changed with resultant ecological succession. Thus, even though many animals were found, they might not be ecologically desirable or valuable to the ecosystem represented by the mangrove community (de Sylva 1972).

Another problem confronting us is that defoliation flights ceased two years prior to our study. Although defoliation may have had severe immediate effects, and even though the mangroves themselves have still not regenerated substantially (as of 1973), perhaps any damage to the equatic habitat and its organisms may not have been permanent. Thus we might be studying an environment that is actually recovering rather than one that has been damaged permanently. Of course, one of the goals of this study has indeed been to determine if any ecological damage is permanent or temporary.

Because collections were made under wartime conditions, it was frequently impossible to collect in areas because of security restrictions imposed by military activity. In some instances, collecting stations could not be repeated in the same area, and hence comparable data were unobtainable. It was also impossible to make nighttime collections because of security reasons. Many aquatic organisms burrow into the mud during the day, so that they cannot be easily caught with nets, and thus they must be captured at night. Further, we were restricted to the use

of a small bottom trawl for the capture of fishes and invertebrates. Larger and more varied types of nets, traps, seines, gill nets, and weirs undoubtedly would have yielded a more representative sample of the fishes of the mangrove ecosystem, but these could not be used for logistic and security reasons. Also, larval fishes and plankton migrate to, and are more numerous at, the surface (where our collections necessarily were made) during the night than they are during the day. Under ideal conditions, then, collections would be made both during night and day.

Of course, the obvious limitation of our study is its short-term nature. Statistical chance dictates that with the spot sampling necessitated by security problems, data will be limited at best.

In the control area, South Vietnamese naval personnel routinely dropped hand grenades into the water to discourage enemy underwater demolition experts (UDT's) or to detonate mines suspected of being planted by enemy UDT's. Hence, the "control area" we selected could not really be considered as being one untouched by human activity.

In both the defoliated Rung-Sat Special Zone and the nondefoliated Vung-Tau Region, the extensive river traffic from large vessels and military patrol craft adds greatly to the already turbid condition of the wacer by stirring up the bottom with propellers and by causing erosion of the stream banks from boat wakes. Resulting siltation and turbidity can reduce productivity in both areas, reducing light penetration and therefore photosynthetic activity of aquatic planktonic and benthonic plants.

Finally, the effects of mangrove defoliation upon the aquatic organisms cannot be separated from other kinds of environmental degradation

(Orians and Pfeiffer 1970). Chlorinated hydrocarbons have been sprayed to control various insect pests throughout Indochina since the end of World War II, and, more recently, organophosphates such as malathion have been added to the list of pesticides. Sewage from the Saigon area and, to ε leaser extent, industrial wastes, accidental spills of oil and other hydrocarbons, and a host of unknown war-related material contaminate the waters flowing through the Rung-Sat Special Zone. The deleterious effects upon the aquatic ecosystem of any pollutant by itself or through its synergistic effect with other pollutants (including defoliants) can only be speculated upon pending necessary laboratory and field studies.

METHODS AND MATERIALS

The primary area of mangrove defoliation in SVN is the Rung-Sat Special Zone (Figure 1), once a region thickly covered by several species of mangrove trees, including <u>Rhizophora</u>, <u>Avicennia</u>, <u>Sonneratia</u>, <u>Bruguiera</u>, and <u>Ceriops</u>. The aquatic environment here is typified by tortuous, deep channels characterized by tidal velocities that may exceed 3 knots (Kopenski 1968). Extensive tidal mixing and stirring occurs (Figure 5), so that the waters are continuously turbid (Table IV). Salt water moves upstream near the river bottom as far as the town of Nha-Be. Above this is a layer of freshwater runoff at all tidal stages that is detectable throughout the Rung-Sat Special Zone.

The control region (the Vung-Tau Region) comprises the Song-Co-May and Song-Mui-Giui (rivers); it is subject to relatively weaker tidal currents (estimated at 1 to 2 knots, maximum) than in the Rung-Sat, and has . .



٠

J.

FIG. 5. Sediment load (% by wt) and current speed (knots) observed in the Saigon, Nha-Be, and Long-Tau Rivers, SVN, Sept. 3-10, 1967, at various tidal stages. U.S. Naval Decanographic Office date (Kopenski 1968). All VND stations are in the defoliated Rung-Sat Special Zone.

15

*

Table IV.

Summary of hydrographic and biological data collected in mangrove region of SVN, 1972-73. The Rung-Sat Special Zone is defoliated; Vung-Tau is nondefoliated (control region). Wet-season collections were made in October and November 1972. Dry-season collections were made in January 1973.

	Rung-Sat wet	<u>Rung-Sat</u> dry	<u>Vung-Tau</u> wet	<u>Vung-Tau</u> dry
Average values for each ar	ea			
Temperature, ^O C	28.8	27.4	29.5	26.8
Oxygen, ppm	4.1	4.4	5.6	6.4
Oxygen saturation, %	67	67	90	98
Salinity, ⁰ /00	11.5	17.0	27.8	30.7
pH	7.5	7.4	8.0	7.9
Turbidity, JTU	62	55	8	4
Number of organisms				
Copepods	127×10^3	117 x 10 ³	273×10^3	114 x 10 ³
Diatoms	106×10^4	456×10^2	511×10^4	229 x 10 ³
Fish eggs	422	2,242	277	1,146
Fish larvae	1,864	10,469	268	274
No. fish families	9	11	15	17
Sponges	1	0	8	6
Corals, etc.	5	6	44	29
Worms	15	24 9+	101	66
Clama, etc.	35	11	104+	28
Crustaceans	783	2,741	1,207	1,712
No. crustacean families	13	18	19	19
Sea stars, etc.	4	2	81	28
Fish	104	215	240	250
No. fish families	17	10	17	16

a shallower average depth. The banks are heavily forested with mangroves, principally <u>Rhizophora</u>, and consequently there seems to be less erosion of the banks, resulting in lower average turbidity values (Tables IV and V). The Vung-Tau Region is not subjected to any sewage or industrial wartes, except for a few villages to the northeast near Phuoc-Le (Ba-Ria) and west of the town of Vung-Tau at Xa-Thang. Throughout its area, the Vung-Tau Region closely resembles the undisturbed mangrove forest of parts of Southwest Florida; thus, in part because some comparable qualitative and quantitative information is available from South Florida, the Vung-Tau Region was considered to be a reasonably useful control area.

Collections were made to obtain representative kinds of animals and plants from many stations over a relatively wide geographical region in both the defoliated and nondefoliated areas. We attempted to characterize the physical environment in each locality in relation to the organisms found there. Lack of time and funds precluded the collection and analysis of the chemical environment or chemical nutrient cycles at each station.

Environmental data were based upon water samples collected at the surface, at a middle depth (half-way to the bottom), and just off the bottom. Water temperature and dissolved oxygen content at each collection station were determined with a battery-operated YSI meter. Water salinity was determined from water samples, which were analyzed at the end of the day using a Brix refractometer. Values for pH were determined in the field using a battery-powered pH meter. The turbidity was determined from water samples collected in the field and was subsequently measured at the

Table V.

Hydrographic and biological data collected in mangrove region of SVN, 1972-73. The Rung-Sat is defoliated; Vung-Tau is nondefoliated (control) region. Wet-season collections were made in October and November 1972. Dry-season collections were made in January 1973. Dashes in columns indicate sample not collected. Maximum sample depth is the depth of the bottom water sample.

RUNG-SAT (defoliated) -- wet season

VND stations	10	16	12	14	32	24	18	20	30	22	28	26	34	Average
Maximum water depth, m	17	15	16	15.	11	25	12	13	11	20	30	3	10	
Maximum sample depth, m	12	11	12	12	8	12	10	12	11	12	12	3	7	
Temperature, ^O C														
Surface sample	28.1	28.8	28.5	28.3	30,0	29.3	28.0	28.9	29.8	29.3	29.6	29.6	29.4	29,0
Middle sample	27.9	28.0	28.2	28.1	30.0	-	28.2	-	28.5	-	29.2	-	29.1	28.6
Bottom sample	28.0	28.0	28.0	28.0	29.8	-	28.2	-	28.9	-	29.2	-	29.1	28.6
Average Oxygen, ppm	28.0	28.3	28.2	28.1	29.9	29.3	28.1	28.9	29.1	29.3	29.3	29.6	2 9. 2	28.8
Surface sample	3.8	4.1	3.9	3.7	4.3	3.6	3.9	3.2	4.6	3.5	5.0	6.0	6.0	4.3
Middle sample	3.7	4.0	3.5	3.5	4.2	3.2	3.4	3.2	4.3	3.5	4.9	-	6.0	4.0
Bottom sample	3.8	3.9	3.4	3.6	4.1	3.3	3.5	3.1	4.4	3.5	4.8	5.6	5.9	4.1
Average % of oxygen saturation	3.8	4.0	3.6	3.6	4.2	3.4	3.6	3.2	4.4	3.5	4.9	5.8	6.0	4.1
Surface sample	59	66	62	58	71	58	61	52	75	56	82	98	97	69
Middle sample	58	63	55	55	69		53	-	68	-	79	-	97	66
Bottom sample	59	61	53	56	67	-	55	-	71	-	77	-	95	66
Average	59	63	57	56	69	58	56	52	71	56	79	9 8	96	67
Surface sample	0.0	0.2	1,1	2.3	5.5	8.1	7.7	12.5	13.1	18.1	17.7	24.1	24.9	10.4
Middle sample	0.0	0.1	1.5	2.5	8.3	12.5	11.3	12.9	12.5	15.3	20.7	-	25.3	10.2
Bottom sample	0.2	0.2	3.3	2.7	17.3	14.5	18,5	13.3	17.9	14.1	23.9	26.7	26.5	14.5
Average	0.1	0.2	2.0	2.5	10.4	11.7	12.5	12.9	14.5	15.8	20.8	25.4	25.6	11.5

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations Maximum water depth, m Maximum sample depth, m	10 17 12	16 15 11	12 16 12	14 15 12	32 11 8	24 25 12	18 12 10	20 13 12	30 11 11	22 26 12	28 30 12	26 3 3	34 10 7	Average
<u>pH</u> Surface sample	6.7	6.7	6.7	7.0	7.4	7.4	7.2	7.5	7.7	7.6	8.0	7.8	8.2	7.4
Middle sample	6.8	7.3	6.9	7.1	7.6	7.6	7.4	7.7	7.8	7.8	8.1	-	8.3	7.5
Bottom sample	6.8	7.2	6.8	7,0	7.7	7.7	7.8	7.6	8.2	7.9	8.1	7.9	8.3	7.6
Average	6.8	7.1	6.8	7.0	7.6	7.6	7.5	7.6	7.9	7.8	8.1	7.9	8.3	7.5
Surface sample	60	49	62	130	20	50	94	220	14	77	13	12	4	62
Plankton, surface ^f														Total
Copepods	658 ⁸	260 ^a	595	347	602 ⁸	742 ^a	538 ^a	137 ^b	862 ⁸	159 ^b	370 ^b	201 ^a	207 ^b	127 ⁰
Diatoms	0	30	686	-	248 ^b	594 ^b	843 ⁸	705 ^b	445 ^b	696 ^b	267 ^C	863 ^b	431 ^e	106 ^đ
Fish eggs	0	0	0	0	0	0	0	0	0	Ó	113	200	109	422
Fish larvae	132	7	22	42	29	37	255	716	7	553	25	5	34	1,864
Number of families	4	1	4	4	4	3	4	3	່ 2	6	3	3	5	9
Benthos ^f Sponges	0	0	0	0	0	0	0	0	0	0	0	0	1e	1
Corals, hydroids, & anemones	0	0	0	0	0	1	0	0	0	2	1	0	1	5
Worms	0	5	0	1	0	0	0	2	0	6	0	1	0	15
Clams, snails, & squids	0	0	0	0	1	6	0	2	4	4	2	12	4	35
^a Number X 10 ¹ ^b Number X 10 ² ^c Number X 10 ³ ^d Number X 10 ⁴ ^e Fragments (considered as one ^f Number of individuals	e indiv	idual)												

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations	10	16	12	14	32	24	18	20	30	22	28	26	34	Total
Maximum water depth, m	17	15	16	15	11	25	12	13	11	20	20	3 ·	10	
Maximum sample depth, m	12	11	12	12	8	12	10	12	11	12	:2	2	7	
Benthos (cont'd.)														
Shrimp & crabs, etc.	21	32	1	6	3	121	388	94	64	32	8	7	6	783
Number of families	2	2	1	4	2	7	6	8	6	6	1	4	3	13
Sea stars, brittlestars, & urchins	0	0	0	0	0	1	0	0	0	1	1	0	1	4
Fishes (benthic)	5	15	1	1	3	16	8	22	2	18	9	4	0	104
Number of fish families	2	5	1	1	3	8	2	6	2	4	6	3	0	17

.

.

-

٠

~

.

`

21

RUNG-SAT (defoliated) -- dry season

VND stations Maximum water depth, m Maximum sample depth, m	66 17 12	64 20 12	62 15 12	60 15 12	80 14 12	74 7 7	72 20 12	68 25 12	70 20 12	78 8 8	76 30 12	Average
<u>Temperature</u> , ^O C Surface sample	28.8	28.1	28.0	27.5	27.8	27.9	28.1	27.2	27.9	27.2	27.3	27.8
Middle sample	28.0	27.7	27.5	27.1	27.3	27.4	27.1	27.0	27.1	27.0	27.0	27.3
Bottom sample	27.9	27.8	27.2	27 .0	27.1	27.2	27.0	27.0	27.0	27.0	27.0	27.2
Average	28.2	27.9	27.6	27.2	27.4	27.5	27.4	27.1	27.3	27.1	27.ì	27.4
Surface sample	3.7	3.8	3.8	5.3	4.1	4.5	4.6	5.1	5.5	5.1	5.3	4.6
Middle sample	3.3	3.3	3.2	4.7	3.9	4.5	4.5	4.6	5,0	4.6	4.9	4.2
Bottom sample	3.0	3.6	3.4	4.6	3.8	4.5	4.5	4.7	5.1	4.4	5.0	4.2
Average	3.3	3.6	3.5	4.9	3.9	4.5	4.5	4.8	5.2	4.7	5.1	4.4
Surface sample	60	59	59	83	64	70	72	78	86	78	82	72
Middle sample	52	52	50	72	60	69	69	71	77	71	75	65
Bottom sample	47	56	52	71	58	69	69	72	78	68	77	65
Average	53	56	54	75	61	69	70	74	80	72	78	67
Salinity, 700 Surface sample	2,3	3,5	6.5	15.9	17.5	14.5	20,9	22.5	24,5	25.7	27.3	16,5
Middle sample	3,1	4.3	6,5	15,7	-	18.9	21.3	23,3	24,9	25 .9	27,3	17.1
Bottom sample	3.5	4.3	6.5	15.5	17,9	20,5	21.3	23.3	26,1	25.9	27.3	17.5
Average	3.0	4.0	6.5	15,7	17.7	18.0	21.2	23.0	25,2	25.8	27,3	17.0
<u>pH</u> Surface sample	6.9	6.0	7.1	7.4	7.1	7.3	7.5	7.7	7.7	7.7	7.8	7.4
Middle sample	6.8	6.9	7.0	7.2	7.2	7.3	7.5	7.6	7.6	7.6	7.8	7.3
Bottom sample	6.8	6.9	7.2	7.5	7.2	7.4	7.5	7.6	7.7	7.6	7.7	7.4
Average	6.8	6.6	7.I	7.4	7.2	7.3	7.5	7.6	7.7	7.6	7.8	7.4

RUNG-SAT (defoliated) -- dry season - (cont[†]d.)

VND stations Maximum water depth, m Maximum sample depth, m	66 17 12	64 20 12	62 15 12	60 15 12	80 14 12	74 7 7	72 20 12	68 25 12	70 20 12	78 8 8	76 30 12	Average
<u>Turbidity, JTU</u> Surface	40	40	120	90	35	22	38	20	50	27	125	55
Plankton, surface ^f												Total
Copepods	691 ^a	144 ^b	507 ^a	598 ^a	217 ^b	434 ^a	239 ^b	752 ²	114 ^b	653 ^a	974 [%]	117 ^c
Diatoms	0	61	206	676	106 ^b	715	981 ⁸	415 ⁸	535 ^a	511 ⁸	891 ⁸	456 ^b
Fish eggs	0	0	0	1	0	0	38	368	325	569	941	2,242
Fish larvae	345	572	1,270	2,212	1,246	530	644	1,429	1,057	470	694	10,469
Number of families	3	4	4	4	4	3	4	7	3	5	5	11
<u>Benthos</u> ^f Sponges	0	0	0	0	0	0	0	0	0	0	0	0
Corals, hydroids, & anemones	0	0	1	1	0	0	0	0	1	0	3	6
Worms	0	0	243+	0	0	1	0	•	0	Q	5	249+
Clams, snails,& squids	0	0	2	0	0	4	0	0	0	0	5	11
Shrimp & crabs, etc.	3	326	337	358	254	121	351	281	307	218	185	2,741
Number of families	2	6	7	3	6	10	2	5	4	6	8	. 18
Sea stars, brittlestars, & urchins	0	0	0	0	0	0	0	0	0	1	1	2
Fishes	1	2	12	1	12	18	10	7	2	137	13	215
Number of families	1	2	3	1	4	7	3	3	2	5	5	10
^a Number X 10 ¹ ^b Number X 10 ² ^C Number X 10 ³ ^d Number X 10 ⁴ ^f Number of individuals												

-

*

22

.

• ,

× 1

.

VUNG-TAU (nondefoliated) -- wet season

e de la companya de l La companya de la comp

23

.

VND stations	42	44	40	48	56	58	50	52	36	38	46	54	Average
Maximum water depth, m	2	3	9	6	3	12	7	6	4	10	5	6	
Maximum sample depth, m Temperature. OC	2	. 3	4	.0	3	8	/	6	4	10	5	6	
Surface sample	29.8	29.8	29.3	29.8	29.9	29.6	29.9	29.8	30.0	29.7	29.8	2 9. 9	29.8
Middle sample	-	-	-	29.2	-	29,0	29.2	29.2	-	-	-	29.8	29.3
Bottom sample	28.8	29.1	29.2	29.1	28.5	29.0	2 9. 8	2 9. 2	2 9. 5	-	29.4	29.8	29.2
Average	29.3	29.5	29.3	29.4	29.2	29.2	29.6	2 9. 4	29.8	29.7	29. 6	29.8	29.5
Surface sample	5.3	4.9	5.3	5.4	5.5	5.7	5.7	5.9	6.5	-	5.4	6.5	5.6
Middle sample	-	-	-	5.3	-	5.8	5.5	5.8	-	-	-	6.5	5.8
Bottom sample	4.9	4.6	4.8	5.3	5.2	5.6	5.2	5.8	5.9	-	5.1	6.3	5.3
Average % of oxygen saturation	5.1	4.8	5.1	5.3	5.4	5.7	5.5	5.8	6.2	-	5.3	6.4	5.6
Surface sample	87.	80	85	89	90	93	93	97	107	-	89	107	92
Middle sample	-	-	-	85	-	94	89	94	-	-	-	107	94
Bottom sample	80	74	77	85	83	90	85	94	97	-	82	103	86
Average	84	77	81	86	87	9 2	89	95	1 0 2	-	86	106	90
Surface sample	21.3	25.7	27.3	27.1	27.3	28.1	28.1	28.3	26.5	28.5	30.1	28.9	27.3
Middle sample	-	-	-	27.5	-	28.1	28.3	28.5	-	28.5	-	29.1	28.3
Bottom sample	26.7	27.1	27.3	27.7	27.9	28.1	28.3	28.7	30.5	28.5	27.5	29.3	28,1
Average pH	24.0	26.4	27.3	27.4	27.6	28.1	28.2	28.5	28.5	28.5	28.8	29.1	27.8
Surface sample	7.6	7.9	7.9	7.6	8.1	8.1	7.7	8.4	8.1	8.3	8.2	8.4	8.0
Middle sample	-	-	-	7.6	-	8.2	7.8	8.4	-	-	-	8.5	8.1
Bottom sample	7.7	7.9	7.9	7.8	8.1	8.2	7.7	8.3	8.1	-	8.2	8.5	8.0
Average	7.7	7.9	7.9	7.7	8.1	8.2	7.7	8.4	8.1	8.3	8.2	8.5	8.0

VUNG-TAU (nondefoliated) -- wet season - (cont'd.)

	VND stations Maximum water depth, m Maximum sample depth, m Turbidity, JTU	42 2 2	44 3 3	40 9 4	48 6 6	56 3 3	58 12 8	50 7 7	52 6 6	36 4 4	38 10 10	46 5 5	54 6 6	Average
	Surface	21	7	7	6	22	8	4	4	3	3	6	4	8
	Plankton, surface f													Total
	Copepods	5 98 ^a	137 ^b	130 ^b	161 ^b	2 04 ^b	386 ^b	405 ^b	416 ^b	405 ^a	222 ^b	222 ^b	342 ^b	273 ^C
	Diatoms	890 ^a	626 ^a	470 ⁸	475 ^b	102 ^C	290 ^e	139 ^d	823 ^C	530 ^e	629 [¢]	407 ^b	124 ^d	511 ^d
	Fish eggs	0	1	0	Ľ	o	3	12	2	180	27	0	51	27 7
	Fish larvae	15 ·	11	6	10	36	12	12	21	113	20	4	8	268
	Number of families	2	4	4	3	5	6	6	4	5	6	2	4	15
	Sponges	0	ıe	1 ^e	0	2 ^e	0	0	2 ^e	0	0	0	2	8
	Corals, hydroids, & anemones	0	1	2	1	0	0	0	· 2	10	9	13	6	44
•	Worms	12	40	0	0	3	0	27	7	1	0	0	11	101
	Clams, snails,& squids	15	0	2	0	11	q	25 1	28	12	0	11	0	104+
	Shrimp & crabs, etc.	26	24	2 1	18	271	77	214	9 3	31	42	124	24	1207
	Number of families	9	8	4	3	4	13	10	7	8	8	11	6	19
;	Sea stars, brittlestars, & urchins	1	32	3	2	2	0	4	28	0	0	3	6	81
	Fishes	55	5	4	14	56	12	23	12	10	17	29	3	240
	Number of families	9	4	3	7	8	6	7	6	6	5	8	1	17
	^a Number X 10 ¹ ^b Number X 10 ² ^C Number X 10 ³ ^d Number X 10 ⁴ ^e Fragments (considered as on	a indi		۰										

.

-

^eFragments (considered as one individual) ^fNumber of individuals

.

κ.

24

.

.

-4

.

VUNG-TAU (nondefoliated) -- dry season

. . . .

	VND stations Maximum water depth, m Maximum sample depth, m	100 8 7	90 17 12	84 6 6	102 16 12	104 13 12	92 16 12	82 16 12	86 14 12	98 8 8	88 17 12	96 17 12	94 25 12	Average
	<u>Temperature, ^OC</u> Surface sample	27.5	27.5	26.8	27.8	27.5	27.9	27.0	27.7	27.3	27.2	27.7	27.3	27.4
	Middle sample	26.8	26.5	26.5	26.6	26.8	26.3	26.3	26.8	26.3	27.0	26.5	26.1	26.5
	Bottom sample	26.3	26.3	26.5	26.5	26.5	26 .2	26.2	26.8	26 .3	26.9	26,0	26.1	26.4
	Average	26.9	26.8	26.6	27.0	26.9	26.8	26.5	27.1	26.6	27.0	26.7	26.5	26.8
	Oxygen, ppm Surface sample	6.8	7.5	5.8	6.8	7.2	7.8	6.1	6.1	6.4	6.9	6.6	6.3	6.7
25	Middle sample	6.5	7.0	5.7	6.4	6.5	6.5	5.9	6.0	6.0	6.8	5.9	6.0	6.3
	Bottom sample	5.9	6.9	5.7	6.3	6.6	6.3	5,8	5.8	6.0	6.8	5.7	5.9	6.1
	Average	6.4	7.1	5.7	6.5	6.8	6.9	5.9	6.0	6.1	6.8	6.1	6.1	б.4
	<u>% of oxygen saturation</u> Surface sample	106	117	89	106	113	122	94	95	98	106	103	9 7	104
	Middle sample	100	108	88	98	100	98	89	92	91	105	91	91	96
	Bottom sample	89	105	88	97	102	95	88	89	91	105	86	89	94
	Average	9 8	107	88	100	105	105	9 0	92	93	105	93	92	98
	<u>Salinity, 700</u> Surface sample	27.7	28.1	30.1	28.9	29.9	27.7	30.7	30.7	29.5	31.3	29.1	30.1	29.5
	Middle sample	29.1	30.7	30,5	30.5	30.7	32,1	30.7	30.7	31.7	31.1	33.5	32.9	31.2
	Bottom sample	30.5	30.9	30.5	31.7	30.7	32.1	30.7	30.7	32.3	31.1	33.5	33.3	31.5
	Average	29.1	29.9	30.4	30.4	30.4	30.6	30.7	30.7	31.2	31.2	32.0	32.1	30.7
	Surface sample	8.0	7.9	7.8	8.1	7.9	8.2	7.5	7.6	8.1	7.7	8.0	8.2	7.9
	Middle sample	8.0	7.6	7.8	8.1	7.9	8.2	7.8	7.9	8.1	7.7	8.0	8.2	7.9
	Bottom sample	8.0	8.1	7.2	8.1	7,9	8.2	7.6	7.7	8.1	7.8	8.0	8.2	7.9
	Average	8.0	7.9	7.6	8.1	7.9	8,2	7.6	7.7	8.1	7.7	8,0	8.2	7.9

VUNG-TAU (nondefoliated) -- dry season - cont'd.)

.

	VND stations Maximum water depth, m Maximum sample depth, m	100 8 7	90 17 12	84 6	102 16 12	104 13 12	92 16 12	82 16 12	86 14 12	98 8 8	88 17 12	96 17	94 25 12	Average
Surface	Turbidity, JTU	3	3	6	4		2	3		3		2	4	4
	Plankton, surface ^f													Total
Copepod	5 ·	100 ^b	362 a	-	353 ^a	574 ^a	106 ^b	262 ^b	789 ^a	12 0 Ъ	137 ^b	879 ^a	115 ^b	114 [°]
Diatoms		572 ^b	304 ^Ъ	-	135 ^b	338 ^b	830 ⁸	285 ^a	165 ^a	235 ^Ъ	377 ^a	299 ^b	244 ^b	229 [°]
Fish eg	gs	109	120	0	15	11	40	4	3	230	2	340	272	1,146
Fish la	rvae	18	8	31	66	21	27	23	14	9	27	13	17	274
Nu	mber of families	3	5	5	7	4	8	4	3	4	2	4	5	17
Sponges	Benthos	0	0	0	0	0	O	4 ^e	0	0	2	0	-	6
Corals,	hydroids, & anemones	4	1	1	2	1	0	5	2	3	2	8	-	29
Worms		0	1	4	0	0	0	11	1	1	48	0	-	66
Clams,	snails, & squids	0	0	3	1	0	1	7	0	1	12	3	-	28
Shrimp 4	£ crabs, etc.	12	40	82	20	50	13	51	125	5	876	438	-	1,712
Nu	mber of families	5	6	10	4	8	5	11	7	2	13	13	-	19
Sea sta	rs, brittlestars, & urchins	0	0	3	0	0	1	3	0	0	18	3	-	28
Fishes		20	72	14	14	8	8	2	0	5	3	104	-	250
Nu	mber of families	5	3	5	7	4	4	2	0	3	2	6	~	16
a N C N C N C N C N C N C N C N C N C N	umber X 10 ¹ umber X 10 ² umber X 10 ³ umber X 10 ⁴ ragments (considered as one umber of individuals	indivi	dual)											

*

¥.

.

2

end of the day using a Hach Jackson turbidimeter. At each station, observations were also made on percentage of cloud cover, wind speed and direction, tidal stage and direction, bottom type, and sightings of fish schools and birds.

Biological collections were made for juvenile and adult fishes and invertebrates using a fine-mesh otter trawl. This 10-ft net was dragged on the bottom, behind the boat (36-ft PBR vessels belonging to the South Vietnamese Navy) for 15 minutes. Simultaneously, two fine-meshed nets of 0.5-m diameter for collecting plankton were towed astern just beneath the surface, one net having a mesh size of 0.5 mm (505 μ), the other having a mesh size of 0.1 mm (110 μ). All samples were immediately preserved in formalin for subsequent examination in the laboratory in Miami.

After the biological samples were returned to the laboratory, they were sorted into general groups (e.g., fishes, crabs, shrimps, worms), and then identified as accurately as possible by specialists familiar with each group. Lack of comparative collections and published literature has hindered identification of most groups of aquatic organisms collected in SVN, even though an intensive search of the literature has been made to determine what published information is required for specimen identification.

Three trips were made to the mangrove regions of SVN. The first, in March 1972, was made in order to select collecting sites, to determine the feasibility of using certain types of equipment, and to ascertain the suitability of the PBRs as collecting vessels. A second

trip was made in October and November 1972, to collect in the Rung Sat Special Zone and the Vung-Tau Region during the monsoon (wet) season, and a third visit was made in January 1973 to the same areas during the dry season. It was believed that different environmental conditions in the estuaries might be found following extensive runoff from rainfall compared to conditions prevalent during the dry season. During the October-November trip, 13 collections were made in the Rung-Sat Special Zone and 12 in the Vung-Tau Region; during the January trip, 11 collections were made in the Rung-Sat Special Zone and 13 were made in the Vung-Tau Region, yielding a total of 49 collections of environmental and biological data during the wet and dry seasons.

RESULTS

Physical Results: Environmental Data

A comparison of the environments of the two areas shows that the average water temperature was approximately the same in both areas (Table IV). As might be expected during the dry season when air temperatures were cooler, water temperatures were proportionately cooler, but nonetheless were similarly comparable in the two regions. It had been expected that, with the loss of shade offered by the mangrove canopy resulting from defoliation in the Rung-Sat Special Zone, solar heating of shallow tidal waters along the river banks might have resulted in higher water temperatures, but the data do not show that this occurred appreciably. In fact, average surface temperature (Table V) was greater in Vung Tau (29.8°C) than in the Rung-Sat (29.0°C), possibly because the average depth is less in the former region, thus permitting solar heating to occur

more quickly. Probably the large volume of water and extensive tidal stirring of the water columns in the Rung-Sat Special Zone quickly mixes any locally heated water with that in the main channels.

Dissolved oxygen, which is a limiting factor in determining the distribution of organisms in the aquatic environment. is statistically lower in the Rung-Sat Special Zone during both wet and dry seasons, and may well be a direct result of defoliation. The large amount of organic plant fibers and rootstalks found suspended in the waters of the Rung-Sat Special Zone presumably removes oxygen from the water due to bacterial decomposition. In spite of considerable water turbulence from tidal mixing, average oxygen saturation values were considerably lower in surface waters of the Rung-Sat Special Zone (69% and 72% in the wet and dry seasons, respectively) than in the Vung-Tau Region (93% and 113% in the wet and dry seasons, respectively). This would suggest that microorganisms are removing oxygen from the Rung-Sat Special Zone, and that high turbidity is inhibiting photosynthesis. While the relatively low values of dissolved oxygen and oxygen saturation in the Rung-Sat Special Zone may not actually kill aquatic animals, in combination with other factors such as the relatively high temperatures and the high turbidity level, low oxygen content is probably a factor affecting production and growth of organisms.

Surface salinity during the wet season, as would be expected, averaged lower in the Rung-Sat Special Zone $(10.4^{\circ}/\circ\circ)$ than in the Vung-Tau Region $(27.3^{\circ}/\circ\circ)$ because of the large influx of fresh water from the Saigon and Dong-Nai Rivers. Even after monsoon rains had subsided, salinities remained lower during the dry season in the Rung-Sat Special Zone $(16.5^{\circ}/\circ\circ)$ than in the Vung-Tau Region $(29.5^{\circ}/\circ\circ)$.

During the wet season in the Rung-Sat Special Zone, there was a distinct surface salinity gradient from zero salinity near Nha-Be, in the north, southward to the coastal zone where the surface salinity was $24.9^{\circ}/\circ$. At all but two stations, salinity was higher at the surface than at the bottom. At three stations, surface salinity was higher than the mid-depth values, possibly reflecting intrusion of water from shallow tidal channels that had become hypersaline from evaporation during periods of relatively high air temperatures. During the dry season, the gradient was similar $(2.3^{\circ}/\circ\circ$ to $27.3^{\circ}/\circ\circ$), though the average surface salinity was higher. At only one station did surface salinity exceed the bottom salinity, thus indicating less hypersaline water entering from shallow channels during the dry season, when air temperatures are lower.

In the Vung-Tau Region, during the wet season, the surface salinity values at one station exceeded that of the bottom sample. Here, with the exception of limited areas of mangrove selectively removed by Vietnamese woodcutters, the mangrove canopy remains relatively intact; hence it would be expected that evaporation of tide pool waters and thus hypersalinity would be less likely to occur. The surface salinity gradients in the Vung-Tau Region were comparable during both the wet $(21.3-30.1^{\circ}/oo)$ and dry $(27.7-30.7^{\circ}/oo)$ seasons; less rainfall was received throughout the year than in the Rung-Sat Special Zone.

Prior to our studies, we had considered that pH might be a limiting factor in preventing germination of mangrove seedlings in the Rung-Sat Special Zone because of changes in soil chemistry following the use of defoliants and napalm, and possibly from extensive craterization. In the Rung Sat Special

Zone, surface pH values in the wet season were 6.7 to 8.2, and in the dry season were 6.9 to 7.9; average values for both seasons were identical (7.4). Values of pH in surface waters of the Vung Tau Region for the wet season were 7.6 to 8.4, and in the dry season were 7.5 to 8.2, with average values of 8.0 and 7.9, respectively. The more acidic surface pH values encountered in the Rung-Sat Special Zone are not considered unusually low for a lower salinity region (Kobayashi 1959). If the defoliation and other war-related activities did affect the soils in the mangrove forest, it is not apparent from our samples taken from the main channels or even in tide pools and shallow channels from the mangrove forests themselves (de Sylva 1972). Of surface samples in the Rung-Sat Special Zone, pH values were lower than mid-depth and bottom samples during the wet season. However, during the dry season, values for surface waters exceeded and equalled samples from mid-depth and bottom waters at all but one station. It is not apparent why pH values of surface waters in the Rung-Sat Special Zone should be so high, especially since salinity values were considerably lower and, especially because pH usually decreases in estuaries as the water becomes less saline. Possibly high pH reflects photosynthetic activity, yet the low percentage of oxygen saturation observed during our daytime collections in the Rung-Sat Special Zone would not indicate high productivity rates.

The last physical measurement examined was that of turbidity, which is an index of water clarity: lower values represent clearer water. For a comparison, the waters of the western shore of southern Biscayne Bay, Florida, a relatively unpolluted, brackish body of water, but revealing the effects of some urbanization, have an average turbidity value of 3.2 JTU

(Jackson Turbidity Units). The average turbidity values for surface waters in the Rung Sat Special Zone for the wet and dry season were 61.9 and 55.2 JTU, respectively; those in the Vung-Tau Region were 7.8 and 3.5 JTU, respectively. This means that the waters of the Rung-Sat Special Zone at comparable seasons were between 8 and 16 times as turbid as the waters of the Vung-Tau Region.

It was not possible to determine what proportion of the turbidity was caused by organic debris or inorganic materials and what was caused by animal and plant plankton. It appeared that most turbidity in the Rung-Sat Special Zone is from organic debris, while that in the Vung-Tau Region is from animal and plant plankton. The striking thing one notices in the Rung-Sat Special Zone is its very high turbidity, with values ranging from a low of 3.5 JTU (near an open, clear bay) to a high of 220 JTU (water resembled coccoa).

Some background information on turbidity in the Rung-Sat Special Zone is available from the U.S. Naval Oceanographic Office (Figure 5). In September 1967, a series of anchor stations in the Saigon, Nha-Be, and Long-Tau Rivers permitted measurements of current speed and the percentage of suspended sediments in the water column of the main channel of the Rung-Sat Special Zone (Kopenski 1968). As the amount of suspended sediment increases, the water clarity decreases and turbidity values correspondingly increase. Examination of the 1967 data shows that the sediment load at that time was quite high, and since these measurements were taken during the period of defoliation of the Rung-Sat, we might assume that the resulting decay of defoliated trees and runoff of debris into the river would have increased the sediment load of the water.
A comparison of the sediment load with current speed, based on the Navy data collected in 1967, shows a good relationship. The faster the current speed, the higher is the concentration of suspended sediments. Because there is a slight lag in concentration of sediments downstream from where each current measurement is made, it is suggested that the tidal current actively stirs up sediments from the bottom and transports them to the surface (upwelling). Further, because of the wide tidal range experienced in the Rung-Sat Special Zone (3-4 m), extensive areas of the eroded shoreline are thus subject to tidal scouring as the tidal current increases. The resulting silt load picked up during tidal scouring would greatly increase the sediment load.

Biological Results

<u>Plankton: Microplankton</u>. Plankton, or the drifting assemblage of small animals and plants, was studied because it offers a good clue to the comparative health of a region. Plankton not only serves as food in the complex aquatic food web for a host of other organisms, but it also comprises larvae of edible shrimps, crabs, clams, oysters, and snails. Because of its large numbers and diversity, plankton can always be found under a variety of conditions, both ideal and adverse. Yet from a careful study of what <u>kinds</u> of plankton are present, it is possible to infer if the plankton is representative of a "healthy" environment, and if the plankton will be eaten by larger organisms to contribute to the economy and success of the aquatic ecosystem. It was thus believed that a thorough analysis of the plankton might give clues, based on knowledge

of planktonic communities from other, similar tropical regions, to the relative health of the mangrove ecosystem in SVN.

In the only comprehensive work attempted on Vietnamese plankton (Shirota 1966), it was explained that studies have exclusively concerned classification of freshwater and marine plankton rather than the ecology of the organisms. Freshwater forms have been studied largely in fish-culture ponds and in lakes and marshes, while the marine plankton has been examined in the relatively oceanic habitat of the Nha-Trang area. A search of the literature and consultation with Vietnamese biologists, including the Directors of Fisheries and of the Cceanographic Institute at Nha-Trang. have confirmed the total lack of material, either samples or data, on estuarine plankton. Inasmuch as there are no data with which to compare the information collected in the Rung-Sat Special Zone and the Vung-Tau Region, we must evaluate the results in terms of reports from somewhat similar areas elsewhere and past experience. The following remarks concern the phytoplankton and zooplankton collected with a 0.1 -mm (110 μ) mesh net at the end of the wet (monsoon) season, October-November 1972, and during the dry season, January 1973.

In the wet season, the brown color and turbidity of the water in the Rung-Sat Special Zone, particularly near the defoliated areas in the region southeast of Nha-Be, led us to believe that the samples contained a large mineral fraction, but this did not prove to be true. Instead, extreme turbidity was directly related to finely-divided debris from higher plants, presumed to be bits of grass and mangrove stems and leaves. This will be discussed later in connection with the abundance of organisms.

The diversity of living plants and animals in both study areas in all except a few samples was far greater than anticipated. Four major kinds of phytoplankton were found--blue-green and green algae, diatoms, and dinoflagellates. These were represented by nine genera (Table VI), but primarily by the diatoms Coscinodiscus spp., which almost invariably dominated the collections. The following ten phyla comprise the zooplankton: Protozoa, Coelenterata, Ctenophora, Nematoda, Mollusca (snail and clam larvae), Annelida (larval and adult worms), Arthropoda (mostly crustaceans), Chaetognatha, Echinodermata (larvae), and Chordata (ascidians and fish larvae). The Arthropoda, numerically the most important among the invertebrate animals, were represented by freshwater cladocerans at stations of 0.0-0.2°/oo salinity (in the Rung-Sat Special Zone) and, at the more saline stations, by larval barnacles, very abundant copepods (approximately 10 species of calanoids, 6 of cyclopoids and 6 of harpacticoids, with large numbers of their larval and juvenile forms), adults and larvae of several shrimp-like animals, and the larvae and juveniles of many kinds of crabs and shrimps (decapods). In all, 52 kinds of invertebrate animals were enumerated (Table VI). With regard to classification, most of the adult planktonic organisms offer fewer problems than the bottom-living invertebrates and the fishes because some species and many genera are cosmopolitan.

A few organisms appear to be indicators of fresh and saline parts of the river, for example the cladocerans <u>Alona</u> sp. and <u>Bosmina</u> sp., which occurred only in fresh water, and the ctenophore <u>Pleurobrachia</u> <u>pileus</u> as well as chaetognaths, which were abundant only at the more saline stations.

Table VI.

List of planktonic organisms collected with 100-µ plankton nets in the Rung-Sat Special Zone and the Vung-Tau Region, SVN, October-November 1972 and January 1973.

PHYTOPLANKTON

ZOOPLANKTON

Cyanophyta, Oscillatoriaceae

Trichodesmium sp.

Chlorophyta, Volvocaceae

Volvox sp.

Chlorophyta, Zygnemataceae

Spirogyra sp.

Chrysophyta, Coscinodiscaceae

Coscinodiscus sp.

Chrysophyta, Chaetoceraceae

Chaetoceros spp.

Chrysophyta, Biddulphiaceae

Triceratium sp.

Chrysophyta, Fragilariaceae

Asterionella sp.

Thalassiothrix sp.

Pyrrophyta, Peridíniaceae

<u>Ceratium</u> sp.

hydrozoan međusae scyphozoan međusae

Protozoa, Radiolaria

Amphilithium sp.

Hippopodius sp.

Ctenophora

Coelenterata

Pleurobrachia pileus

Aschelminthes, Nematoda

Mollusca

gastropod veligers

pelecypod veligers

Annelida

<u>Sagitella</u> sp.

polychaete larvae

Arthropoda, Crustacea

Cladocera

<u>Alona</u> sp. <u>Bosmina</u> sp.

Copepoda, Calanoida

Acartia sp. 1

Acartia sp. 2

Eucalanus sp.

Euchaeta sp.

Labidocera sp.

Paracalanus sp.

Copepoda, Cyclopoida

Corycaeus sp.

Cyclops sp.

Oithona sp.

Oncaea sp.

cyclopoid sp. 1

cyclopoid sp. 2

Temora sp.

Pseudodiaptomus sp.

Centropages sp. 1

Centropages sp. 2

Copepoda, Harpacticoida

Canthocamptus sp.

<u>Clytemnestra</u> sp.

Euterpe sp.

Euterpina sp.

Laophonte sp.

Microsetella sp.

Cirripedia

nauplii

metanauplii

cypris larvae

...

Cunacea

Isopoda

cryptoniscus larvae

Amphipoda

<u>Vibilia</u> sp.

Decapoda

Lucifer typus

Lucifer zoea and mysis

caridean loea and mysis brachyuran loea and mysis

biddiydidi. Socia and i

Stomatopoda

Squilla larvae

Chaetognatha

Sagitta enflata

S. neglecta

<u>S. friderici</u>

Echinodermata

ophiopluteus larvae

juvenile asteroids

Chordata

ascidian larvae

Oikopleura sp.

fish larvae

To evaluate the distribution and abundance of plankton in the estuary, copepods, crab larvae, chaetognaths, and diatoms were selected as guides. At all stations except in fresh water, total numbers of copepods of all species and their developmental stages and of the diatom <u>Coscinodiscus</u> were relatively high. However, these numbers were distinctly lower at stations near defoliated areas, where the salinity did not exceed $20^{\circ}/\circ\circ$, than at stations with higher salinities near nondefoliated mangrove stands. Probably in part owing to high levels of nutrients during the wet*season, the majority of these most abundant groups were then far more numerous than in the dry season. Only mollusk and crab larvae and the copepod genus <u>Acartia</u> occurred in significantly higher numbers in the dry season. These data are summarized in Table VII, in which average numbers of organisms/m³ at all stations combined are compared.

In wet and dry seasons, diatoms showed strong correlation with low turbidity and high salinity. Although they were present over a wide range of salinity (6.5-32.0°/00), the greatest numbers in both seasons were found in the most saline, least turbid waters. Copepods, however, were abundant in almost all parts of both study areas. In the wet season, 80 percent occurred in waters where the turbidity measured 3-20 JTU, and 20 percent occurred in waters of 21-220 JTU, but in the dry season the percentages were approximately equal, owing to the abundance of certain kinds (the calanoid <u>Acartia</u> and species of cyclopoids) that are particularly tolerant of low salinity. It appears that these are organisms that contribute to the high turbidity of the less saline waters.

Chaetognaths, perhaps in response to the abundance of copepods on

Table VII.

Average number of organisms collected in a cubic meter of water in wet and dry seasons, in the Rung-Sat Special Zone and the Vung-Tau Region, SVN, October-November 1972 and January 1973.

Average No./ m ³ , All Stations Combined											
GROUP	WET	DRY									
Coscinodiscus spp.	246,693	12,487									
Total Copepoda [®]	15,975	10,507									
Calanoid Copepoda	6,916	5,132									
Calanoida, <u>Acartia</u>	799	1,209									
Cyclopoid Copepoda	7,149	4,031									
Harpacticoid Copepoda	204	281									
Brachyuran larvae	34	1,299									
Gastropod veliger larvae	152	1,896									
Pelecypod larvae	301	444									

^aThis figure includes adults, juveniles, and identifiable larvae as well as nauplii which could not be assigned to order. which they feed, were often common in areas of relatively low salinity $(15-20^{\circ}/\circ\circ)$.

Crab larvae were rare during the wet season when they were strongly associated with low turbidity and high salinity. Conversely, they were among the most numerous organisms in the dry season and 94 percent occurred. in waters of 21-125 JTU rather than in less turbid waters. Probably a breeding period accounts for the large difference in total numbers, and the presence of different species may well explain the change in environmental preference.

The actual percentage distribution of these key groups in wet and dry seasons, arranged both in order of increasing turbidity and increasing salinity, are shown in Tables VIII and IX.

Larvae of snails and bivalve mollusks (gastropod and pelecypod veligers) occurred in considerable numbers over most of the range of conditions, and often they were abundant in the Rung-Sat region. For example, 6753 snail larvae/m³ of water filtered were collected at a station where the salinity was 6.5°/oo and the turbidity 120 JTU; 1388 bivalve larvae/m³ were taken in water measuring 4.0°/oo and 40 JTU. Snail larvae were lacking at stations of highest turbidity (Station Nos. 15 and 21) during the wet season but were very numerous during the dry season, as shown in the example just given. Clam larvae, never as numerous, were most abundant in the Vung-Tau Region during the wet season but occurred through a broader range of conditions in dry season. Because several species of both groups are involved, each with its own breeding behavior, and because the larvae cannot now be identified to species, it cannot be

Table VIII.

Percentages of groups of organisms arranged in order of increasing salinity at stations sampled during wet and dry seasons in the Rung-Sat Special Zone and Vung-Tau Region, SVN, October-November 1972 and January 1973. Figures for salinity ($^{\circ}/_{\circ\circ}$) and percentages of samples are expressed as X 10.

VET_SEASON_DATAª

DRY_SEASON_DATA

	a . +			1	• • • • • •	STA	SAL	COSC	COPE	BRAC	CHAE
STA	SAL	cosc	COPE	BRAC	CHAE	6 7	30	0	30	1	0
11	1	0	16	2	0	65	40	0	62	1	Ō
17	2	0	7	7	0	63	65	1	22	3	ō
13	20	0	1	10	0	61	157	2	26	ģ	ō
15	25	0	1	25	0	81	177	39	94	112	53
33	104	4	15	13	0	75	180	3	19	R	11
25	117	10	19	14	1	73	212	36	103	129	187
19	125	1	13	20	10	69	230	15	33	<u> </u>	30
21	129	11	34	5	· 7	' 71	252	19	цq	467	53
31	145	7	22	32	0	79	258	19	28	63	
23	158	11	40	178	87	77	273	32	<u>ь</u> р	140	<u>н</u> о
29	208	43	93	42	41	101	201	202	Z - L- Z	2 4 3	101
4 3	240	1	15	0	0	91	291	111	16		20
27	254	14	5	10	2	. 103	200	+++	15	5	20
35	256	70	52	116	50	105	304	100	10		
45	264	1	34	0	8	. 103	304	20	2.3		107
41	273	1	33	7	5	. 30	300	10	40		107
49	274	8	40	17	16	67	207	10	213	2	21
57	276	17	51	1 19	66	.07	307	0	54	4	ن برع
59	281	47	97	8	26	33	342	4 1	52	2	04
51	282	225	102	32	235	09	313	100	23	~	20
37	285	133	104	50	73	5/	320	109	30	0	33
39	285	86	10	8	22	95	371	89	20	v	90
53	285	102	56	182	177						
47	288	7	56	2	52						
55	291	200	86	201	121						

^A Abbreviations: STA = station, SAL = salinity, COSC = <u>Coscinodiscus</u> (diatom), COPE = total Copepoda, including larval and juvenile forms, BRAC = Brachyuran (crab) larvae, CHAE = Chaetognatha.

Percentage of groups of organisms arranged in order of increasing turbidity at stations sampled during wet and dry seasons in the Rung-Sat Special Zone and Vung-Tau Region, SVN, October-November 1972 and January 1973. Turbidity is measured in JTU; percentages of samples are expressed as X 10.

NET_SEASON_DATA^a

DRY_SEASON_DATA

STA	TURB	COSC	COPE	BRAC	CHAE	STA	TURB	COSC	COPE	BRAC	CHAE
35	3	70	52	116	50	93	2	30	46	4	107
37	3	133	104	50	73	97	2	109	38	0	33
39	3	86	10	8	22	83	3	10	113	1	27
53	3	102	56	182	177	91	3	111	16	0	22
51	4	225	102	32	235	99	3	85	52	1	64
55	4	200	86	201	121	101	3	208	43	3	191
49	6	8	40	17	16	105	Э	123	25	1	11
47	6	7	56	2	52	87	4	6	34	2	3
45	7	1	34	0	8	95	4	89	50	0	90
41	7	1	33	7	5	103	4	49	15	2	30
59	8	47	97	8	26	89	5	14	• 59	2	26
27	12	14	5	10	2	69	20	15	33	41	30
29	13	43	93	42	41	75	22	3	19	8	11
31	14	7	22	32	0	. 79	27	19	28	63	24
33	20	4	15	13	0	81	35	39	94	112	53
43	21	1	15	0	0	73	38	36	103	129	187
57	22	17	51	19	66	65	40	0	62	1	0
17	49	0	7	7	0	67	40	0	30	1	0
25	50	10	19	14	1	71	50	19	49	467	53
11	60	0	16	2	0	61	90	2	26	9	0
13	62	0	1	10	0	63	120	1	22	3	0
23	77	11	40	178	87	77	125	32	42	149	40
19	94	1	13	20	10						
15	130	0	1	25	0						
21	220	11	34	5	7						

^a Abbreviations: STA = station, TURB = turbidity, COSC = <u>Coscinodiscus</u> (diatom), COPE = total Copepoda, including larval and juvenile forms, BRAC = Brachyuran (crab) larvae, CHAE = Chaetognatha. stated that edible species are reproducing, only that such numbers and diversity of larval life suggest a viable system.

Very little evidence of seasonal restriction was found. Stomatopod larvae were caught only during the dry season and at rather low salinities, ranging from $17.7^{\circ}/_{\circ\circ}$, where they were most numerous, to $27.3^{\circ}/_{\circ\circ}$. Also, echinoderm larvae, very rare in the wet season, were comparatively widespread in the dry season, occurring in low numbers at 16 of the 22 stations.

In summary, both study areas are rich both in variety of planktonic organisms and in numbers of individuals. These values increase, as one would expect in the tropics, with an increase in salinity. The other environmental parameter measured that appears to have influenced diversity and density of plankton is turbidity. (As far as invertebrate plankton is concerned, the range of temperature is miniscule, oxygen was not at limiting levels, and the relationship, if any, of pH was obscure.) At stations near defoliated areas, turbidity was very high, reaching 220 JTU. Although the average numbers of the most abundant organisms per cubic meter of water in the Rung-Sat Special Zone were far lower than at stations having higher salinity values, the average volume of organic material was much higher in the Rung-Sat Special Zone. This fact results mainly from the large volume of organic debris, consisting principally of decayed fragments of higher plants. These persistent bits of cellulose are kept in suspension by tidal action and the propellers and wakes of river traffic. It is reasonable to infer that the resultant reduction of light in the water column directly affects productivity in this portion of the Rung-Sat Special Zone. In contrast, the high

volume of organic material in the Vung-Tau Region reflects the abundance of phytoplankton and zooplankton, rather than detritus. The huge numbers of <u>Coscinodiscus</u> as well as copepods and larval forms indicate a healthy and richly productive environment in the Vung-Tau Region.

Plankton: Fish Eggs and Larvae. Larval fishes are important indicators of water quality because adults can only spawn in waters that are reasonably free from contaminants (Hynes 1971). Eggs thus spawned either float at or near the surface, or sink to or near the bottom, where they may become attached to vegetation. The eggs must be subjected to optimum environmental conditions if they are to survive to hatch and if the developing larval stages are subsequently to grow into the adult stage (Hempel 1965). Hence, polluted environments generally do not permit the production of large numbers of eggs and larvae by many species of fish (de Sylva 1970). However, occasionally pollution will result in nutrient fertilization of the aquatic environment so that such habitata may be actually favorable to some fish species (Mansueti 1962). When this occurs, usually the number of different kinds of species able to spawn is drastically reduced, and one or several tolerant forms survives to occupy the living space vacated by the more sensitive organisms that have not survived. This results in a few tolerant species producing large numbers of eggs; consequently a large fish population with a great number of only a few species evolves. In contrast, a healthy environment comprises a large number of different species, each producing a moderate number of eggs. Thus, the relative "health" of an area also can be examined by comparing the species diversity, or the number of different

kinds of organisms present, rather than assessing the total number of organisms present.

It is virtually impossible to identify preserved fish eggs. However, the total numbers of fish eggs were enumerated because the presence of fish eggs usually indicates that adult fishes must have spawned nearby, and that, presumably, this necessarily occurs in a relatively healthy environment.

Fish eggs were 1.5 to 2 times as numerous in the Rung-Sat Special Zone during the wet and dry seasons, respectively, as they were in the Vung-Tau Region (Table IV). In the Rung-Sat Special Zone, eggs were taken at 3 of 13 stations (23%) during the wet season and at 6 of 11 stations (55%) during the dry season. In comparison, in the Vung-Tau Region, eggs occurred at 8 of 12 stations (67%) during the wet season and at 11 of 12 stations (92%) during the dry season. The explanation may be that a few very fecund individuals spawned a large number of eggs in a small area, a situation typical of stressed environments.

In the Rung-Sat, low values of pH, oxygen, or both may be factors limiting the spawning and survival of eggs. For example, during the wet season, eggs were taken only when the pH exceeded 7.8, and during the dry season only when the pH exceeded 7.4 (Table V). However, since surface pH values below 7.6 were not encountered in the Vung-Tau samples, comparisons are not possible. A more meaningful relationship appears to exist between the presence of fish eggs and the percent of oxygen saturation in surface waters. In the Rung-Sat Special Zone, no eggs were taken during the dry season when the oxygen saturation was below 72 percent, and during the wet

season none occurred when the saturation was below 82 percent. In the Vung-Tau Region, eggs were usually absent when the oxygen saturation fell below 90 percent.

Larval fishes in the Rung-Sat Special Zone were from nearly 7 times to 38 times as abundant as in the Vung-Tau Region during the wet and dry seasons, respectively. Larval fishes occurred at all stations during both seasons and in each study area.

Identification of larval fishes in the estuarine region of Vietnam is extremely difficult because of the lack of studies there, and because literature on larvae is virtually nonexistent. Nonetheless, it was possible to identify many specimens to the family level (Table X). In the Rung-Sat Special Zone, 9 and 11 different families of fishes were represented in larval fish collections during the wet and dry seasons, respectively, while 15 and 17 families occurred in Vung-Tau collections during the same periods.

Within the Rung-Sat Special Zone, the Gobiidae accounted for 93 percent of larvae identified during the wet season and 98 percent of those collected during the dry season. In the Vung-Tau Region, this family was also predominant, but represented only 57 percent of collections during the wet season and 47 percent of the fish families obtained during the dry season. In studies of southern Biscayne Bay, Florida, gobies are indicators of stressed environments (de Sylva 1970); where pollution or alteration of the ecosystem has occurred, this family usually dominates the fish plankton. Thus, the preponderance of Gobiidae in the Rung-Sat Special Zone would indicate a stressed environment, in contrast to a more rich diversification of organisms in the Vung-Tau Region.

Table X.

.

Larval fish families collected with 0.5-m (505-µ mesh) plankton net in mangrove region of SVN, 1972-73. Numbers in columns are numbers of specimens. The Rung-Sat is defoliated; Vung-Tau is nondefoliated (control) region. Wet season collections were made in October and November 1972. Dry season collections were made in January 1973.

			RUNG	-SAT (defoli	ated)	wet	seaso	n					
VND stations	10	16	17	14	32	18	24	20	30	- 28	22	26	34	Total
Surface salinity, ⁰ /00	0,0	0.2	1.1	2.3	5.5	7.7	8.1	12.5	13.1	17.7	18.1	24.1	24.9	
Fish families (herrings)	0	0	1	1	5	1	1	14	0	0	0	2	19	44
e (anchovies)	1	0	4	4	0	5	1	2	0	0	4	0	0	21
lae (pipefishes & seahorse	es) 1	0	0	0	0	0	0	0	0	0	1	0	0	2
nidae (bigeyes)	0	0	0	0	0	0	0	0	2	0	0	0	0	2
le (jacks)	0	0	0	0	0	0	0	0	0	0	0	0	1	1
porgi es)	0	0	Ð	0	0	0	0	0	0	1	2	1	2	6
e (blennies)	3	0	0	2	1	1	0	0	0	14	3	2	0	26
(naked blennies)	0	0	1	o	1	0	0	0	0	0	1	0	8	11
gobies)	16	1	10	30	20	. 84	32	698	2	7	513	0	2	1415
able specimens	18	6	б	2	0	2	2	0	1	1	18	0	0	56
able (damaged) specimens	93	0	0	3	2	162	1	2	2	2	11	0	2	280
imens	132	7	22	42	29	255	37	716	7	25	553	5	34	1864
	0	0	0	0	0	0	0	0	0	113	0	200	109	422
	VND stations Surface salinity, ⁰ /00 <u>Fish families</u> (herrings) e (anchovies) ae (pipefishes & seahorse idae (bigeyes) he (jacks) porgies) (blennies) (naked blennies) gobies) able specimens able (damaged) specimens imens	VND stations10Surface salinity, °/000.0Fish families (herrings)0(herrings)0e (anchovies)1ae (pipefishes & seahorses)1idae (bigeyes)0e (jacks)0porgies)0(blennies)3(naked blennies)0gobies)16able specimens18able (damaged) specimens93cimens13200	VND stations1016Surface salinity, °/000.00.2Fish families (herrings)00e (anchovies)10ae (pipefishes & seahorses)10idae (bigeyes)00idae (bigeyes)00e (jacks)00porgies)00(hernies)30(naked blennies)00gobies)161able specimens1327imens1327000	ND stations 10 16 17 Surface salinity, °/00 0.0 0.2 1.1 Pish families (herrings) 0 0 1 (herrings) 0 0 1 e (anchovies) 1 0 4 ae (pipefishes & seahorses) 1 0 0 idae (bigeyes) 0 0 0 idae (bigeyes) 0 0 0 porgies) 0 0 0 (blennies) 3 0 0 (gobies) 16 1 10 able specimens 18 6 6 able (damaged) specimens 93 0 0 imens 132 7 22 0 0 0 0	RUNG-SAT (VND stations 10 16 17 14 Surface salinity, ⁰ /00 0.0 0.2 1.1 2.3 Fish families (herrings) 0 0 1 1 e (anchovies) 1 0 4 4 ae (pipefishes & seahorses) 1 0 0 0 idae (bigeyes) 0 0 0 0 idae (bigeyes) 0 0 0 0 idae (bigeyes) 0 0 0 0 idae (biennies) 0 0 0 0 igobies) 16 1 10 30 igobies) 16 1 10 30 able specimens 18 6 2 2 imens 132 7 22 42	RUNG-SAT (defolied VND stations 10 16 17 14 32 Surface salinity, 0/00 0.0 0.2 1.1 2.3 5.5 Pish families (herrings) 0 0 1 1 5 e (anchovies) 1 0 4 4 0 ae (pipefishes & seahorses) 1 0 0 0 0 idae (bigeyes) 0 0 0 0 0 0 idae (bigeyes) 0 0 0 0 0 0 porgies) 0 0 0 0 0 0 0 (blennies) 3 0 2 1 1 30 20 gobies) 16 1 10 30 20 able specimens 18 6 2 0 able (damaged) specimens 93 0 3 2 2 2 2 0 0 0 0 0 3 2 2 2 2 0<	VND stations 10 16 17 14 32 18 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 Fish families (herrings) 0 0 1 1 5 1 e (anchovies) 1 0 4 4 0 5 ae (pipefishes & seahorses) 1 0 0 0 0 0 idae (bigeyes) 0 0 0 0 0 0 0 porgies) 0 0 0 0 0 0 0 0 gobies) 16 1 10 30 20 84 able specimens 18 6 2 0 2 able (damaged) specimens 93 0 0 3 2 162 imens 132 7 22 42 29 255	RUNG-SAT (defoliated) wet VND stations 10 16 17 14 32 18 24 Surface salinity, ⁰ /00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 <u>Fish families</u> (herrings) 0 0 1 1 5 1 1 e (anchovies) 1 0 4 4 0 5 1 ae (pipefishes & seahorses) 1 0 4 4 0 5 1 ae (pipefishes & seahorses) 1 0 0 0 0 0 0 0 uidae (bigeyes) 0 0 0 0 0 0 0 0 porgies) 0 0 0 0 0 0 0 0 (blennies) 3 0 2 1 1 0 gobies) 16 1 10 30 20 84 32 able (damaged) specimens 93 0 0 3 2 162 1 <	RUNG-SAT (defoliated) wet seaso VND stations 10 16 17 14 32 18 24 20 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 Pish families (herrings) 0 0 1 1 5 1 1 14 e (anchovies) 1 0 4 4 0 5 1 2 ae (pipefishes & seahorses) 1 0 4 4 0 0 0 0 iddae (bigeyes) 0 0 0 0 0 0 0 0 0 porgies) 0 0 0 0 0 0 0 0 0 (blennies) 3 0 0 1 0 0 0 0 0 0 gobies) 16 1 10 30 20 84 32 698 able (damaged) specimens 93 0 0 3 2 16 1 2	RUNG-SAT (defoliated) wet season VND stations 10 16 17 14 32 18 24 20 30 Surface salinity, ⁰ /00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 <u>Fish families</u> (herrings) 0 0 1 1 5 1 1 14 0 e (anchovies) 1 0 4 4 0 5 1 2 0 ae (pipefishes & seahorses) 1 0 4 4 0 5 1 2 0 oidae (bigeyes) 0	NND stations 10 16 17 14 32 18 24 20 30 28 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 Fish families (herrings) 0 0 1 1 5 1 1 14 0 0 e (anchovies) 1 0 4 4 0 5 1 2 0 0 ae (pipefishes & seahorses) 1 0	NNG-SAT (defoliated) wet season VND stations 10 16 17 14 32 18 24 20 30 28 22 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 Pish families (herrings) 0 0.1 1 5.5 7.7 8.1 12.5 13.1 17.7 18.1 Pish families (herrings) 0 0.1 1 5 1 1 14 0 0 0 e (anchovies) 1 0 </td <td>ND stations 10 16 17 14 32 18 24 20 30 28 22 26 Surface salinity, ⁰/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 24.1 Pish families (herrings) 0 0 1 1 5 1 1 14 0 0 0 2 e (anchovies) 1 0 4 4 0 5 1 1 14 0 0 4 0 ae (pipefishes & seahorses) 1 0<td>NNDC-SAT (defoliated) wer season VND stations 10 16 17 14 32 18 24 20 30 28 22 26 34 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 24.1 24.9 Pish families (herrings) 0 0.0 1 1 55 1 1 14 0 0 0 2 19 e (anchovies) 1 0 0 1 15 1 1 14 0</td></td>	ND stations 10 16 17 14 32 18 24 20 30 28 22 26 Surface salinity, ⁰ /00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 24.1 Pish families (herrings) 0 0 1 1 5 1 1 14 0 0 0 2 e (anchovies) 1 0 4 4 0 5 1 1 14 0 0 4 0 ae (pipefishes & seahorses) 1 0 <td>NNDC-SAT (defoliated) wer season VND stations 10 16 17 14 32 18 24 20 30 28 22 26 34 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 24.1 24.9 Pish families (herrings) 0 0.0 1 1 55 1 1 14 0 0 0 2 19 e (anchovies) 1 0 0 1 15 1 1 14 0</td>	NNDC-SAT (defoliated) wer season VND stations 10 16 17 14 32 18 24 20 30 28 22 26 34 Surface salinity, °/00 0.0 0.2 1.1 2.3 5.5 7.7 8.1 12.5 13.1 17.7 18.1 24.1 24.9 Pish families (herrings) 0 0.0 1 1 55 1 1 14 0 0 0 2 19 e (anchovies) 1 0 0 1 15 1 1 14 0

RUNG-SAT (defoliated) -- dry season

VND stations	66	64	62	74 ·	60	80	72	68	70	78	76	Totel
Surface salinity, ⁰ /00	2.3	3.5	6.5	14.5	15.9	17.5	20.9	22.5	24.5	25 7	27.3	
<u>Fish</u> <u>families</u> Clupeidae (herrings)	45	30	8	0	5	1	1	9	0	1	3	103
Engraulidae (anchovies)	0	C	6	0	1	0	0	3	1	1	12	24
Syngnathidae (pipefishes & seahorse	s) 0	0	0	3	2	7	3	1	5	1	3	25
Atherinidae (silversieds)?	5	2	2	0	0	1	1	1	0	0	0	12
Sillaginidae (whitings)	0	0	0	0	0	0	0	1	0	0	0	1
Blenniidae (blennies)	0	0	0	0	0	0	0	0	0	1	0	1
?Eleotridae (sleepers)	0	0	0	1	0	0	0	0	0	0	0	1
Gobiidae (gobies)	293	502	1253	525	21 96	1206	628	1 3 44	1043	421	659	10,070
Heterosomata (flounders & soles)	0	3	0	0	0	0	0	0	0	0	0	3
Tetraodontidae (puffers)	0	0	0	0.	0	0	0	1	0	0	0	1
Diodontidae (porcupine fishes)	0	0	0	0	0	0	0	0	0	0	1	1
Unidentifiable specimens	2	18	1	0	8	10	10	37	8	43	16	153
Unidentifiable (damaged) specimens	0	17	0	1	0	21	1	32	0	2	0	74
Total specimens	345	572	1270	530	221 2	1246	644	1429	1057	470	694	10,469
Fish eggs	0	0	0	0	1	0	. 38	368	325	569	941	2,242

.

.

.

.

.

.

.

-

.

• .

. .

VUNG-TAU (nondefoliated)	wet	season	
--------------------------	-----	--------	--

VND stations	42	44	36	48	40	56	50	58	52	:8	54	46	Total
Surface salinity, ⁰ /00	21.3	25.7	26.5	27.1	27.3	27.3	28.1	28.1	28.3	28.5	28.9	30.1	
Fish families Clupeidae (herrings)	0	1	1	0	0	1	0	0	0	0	0	0	3
Engraulidae (anchovies)	0	0	0	0	0	0	l	0	0	1	0	Û	2
Syngnathidae (pipefishes & seahorses)	0	1	0	0	0	0	0	1	1	3	1	0	7
Mugilidae (mullets)	0	0	0	0	0	0	1	0	0	0	l	0	2
Atherinidae (silversides)	0	0	0	1	0	0	0	0	0	0	0	0	1
Centropomidae (snooks)	0	0	0	0	0	23	0	0	6	0	0	0	29
Sillaginidae (whitings)	7	0	0	0	0	0	0	0	0	0	0	0	7
?Carangidae (jacks)	0	0	0	0	1	0	0	0	0	0	0	0	1
Sciaenidae (croakers)	0	0	2	0	0	0	0	1	0	1	0	0	4
Sparidae (porgies)	0	0	1	0	0	1	1	0	0	0	0	0	3
Scaridae (wrasses)	0	0	0	0	0	0	0	1	0	0	0	0	1
Blenniidae (blennies)	0	0	0	0	1	0	1	0	0	0	0	0	2
?Clinidae (naked blennies)	0	3	3	3	1	4	7	1	8	4	0	3	37
Gobiidae (gobies)	8	4	93	1	3	6	1	5	2	8	2	1	134
Heterosomata (flounders & soles)	0	0	0	0	0	0	0	1	0	1	1	0	3
Unidentifiable specimens	0	2	7	2	0	1	0	2	1	2	3	0	20
Unidentifiable (damaged) specimens	0	0	6	3	0	0	0	0	3	0	0	0	12
Total specimens	15	11	113	10	5	36	12	12	21	20	8	4	268
Fish eggs	0	1	180	1	Ũ	0	12	3	2	27	51	0	277

	VUNG-TAU (nondefoliated) dry season												
VND stations	92	100	90	102	96	9 8	104	84	94	82	86	88	Total
Surface salinity, ⁰ /00	27.7	27.7	28.1	28.9	29.1	29.5	29.9	30. 1	30.1	31.7	30,7	31 .3	
<u>Fish families</u> Clupeidae (herrings)	2	0	2	6	4	· 0	0	0	2	0	0	0	16
Engraulidae (anchovies)	1	0	1	5	0	0	1	0	0	0	0	0	٤
Syngnathidae (pipefishes & seahorses)	0	0	0	0	0	0	0	4	0	1	0	1	ć
Mugilidae (mullets)	0	0	· 0	0	0	2	1	0	0	0	0	0	3
Atherinidae (silversides)	0	0	0	0	0	0	0	0	0	1	0	0	1
Centropomidae (snooks)	0	0	1	0	0	0	0	0	0	0	1	0	2
Sillaginidae (whitings)	1	0	0	0	0	0	1	0	2	0	0	Ð	4
?Carangidae (jacks)	1	0	0	3	0	0	0	0	0	0	0	0	4
Pomadasyidae (grunts)	0	0	0	. 3	0	0	0	0	0	2	0	0	5
Sciaenidae (croakers)	3	3	0	1	1	1	0	2	2	0	2	0	15
Sparidae (porgies)	3	2	0	0	0	2	0	1	4	0	0	0	12
Scaridae (wrasses)	1	0	0	Q	0	0	0	0	0	0	0	0	1
Blenniidae (blennies)	0	0	0	0	0	0	0	1	0	0	2	0	3
Clinidae (naked blennies)	0	0	1	0	1	0	0	0	0	0	0	0	2
Gobiidae (gobies)	10	12	0 .	27	0	2	1	9	2	5	0	9	77
Platycephalidae (flatheads)	0	0	3	0	0	0	0	0	0	0	0	0	2

50

.

VUNG-TAU (nondefoliated) dry season (cont'd.)													
VND stations	92	100	90	102	9 6	98	104	84	94	82	86	88	Total
Surface salinity, 0/00	27.7	27.7	28.1	28.9	2 9. 1	29.5	29.9	30.1	30.1	30.7	30.7	31.3	
<u>Fish families</u> Heterosomata (flounders & soles)	0	0	0	1	1	0	0	0	0	0	0	0	2
Unknown specimens	5	1	0	19	4	0	17	14	5	13	6	17	101
Unidentifiable (damaged) specimens	0	0	0	1	2	2	0	0	0	1	3	0	9
Total specimens	~~~27.	18	8	66	13	9	21	31	17	23	14	27	274
Fish eggs	40	109	120	15	340	230	11	0	272	4	3	2	1146

Another measure of richness is given in the variety index (Odum 1971):

d = S \sqrt{N}

where d = the diversity index, S = number of species (families in this case), and N = number of individuals.

This index was calculated for the larval fish families listed in Table X, and the values are presented in Table XI. At nearly every station in the Rung-Sat Special Zone, the values are lower than those obtained in the Vung-Tau Region; the weighted index for all stations in the Rung-Sat Special Zone is somewhat higher during the wet season, but the diversity index for the Vung-Tau Region is higher during the dry season. This greater variety of families is, however, partly due to the more saline regime characteristic of the Vung-Tau Region. Nonetheless, collections made in the Rung-Sat Special Zone during the dry season, when salinities are higher (Table IV), still do not reflect the increase in species diversity one might find in regimes of higher salinity.

Because larval fishes can detect visually and thus avoid an oncoming plankton net in clear water, such as is found in South Florida, larger nets are desirable. The use of 1-m nets would obviate this avoidance reaction but 1-m nets could not be used in Vietnam because the large volume of accumulated floating debris in the water would make the net impossible to retrieve. Another factor that should be taken into consideration is that the clearer water in the Vung-Tau Region may permit larval fishes to escape the oncoming net, hence resulting in a lower catch/m³. Conversely, larvae living in the highly turbid waters of

Table XI.

π.

.

Diversity indices of larval fish families from surface plankton tows in estuaries of SVN. Stations with comparable surface salinity values are in adjacent columns. Stations are arranged in order of increasing surface salinity.

	RUNG-SAT	SPECIAL ZONE		VUNG-TAU REGION									
Wet Seas	<u>nc</u>	Dry Season	<u>n</u>	<u>Wet</u> Seas	on	Dry Seaso	<u>20</u>						
VND stations	<u>index</u>	VND stations	<u>index</u>	VND stations	index	VND stations	index						
10	0.87	66	0.22	42	0.52	92	1.73						
16	1.00	64	0.21	44	1.67	100	0,94						
12	1.00	62	0.14	36	0,60	90	1.77						
14	0.66	74	0,13	48	1.79	102	0.99						
32	0.78	60	0,14	40	1.63	96	1.51						
18	0.35	80	0.14	56	1.01	98	1.51						
24	0.52	72	0.20	50	1.73	104 ·	1.07						
20	0.11	68	0.21	58	2.22	84	1.08						
30	1.00	70	0.12	52	1,21	94	1.46						
28	0.64	78	0.28	38	1.65	82	1.07						
22	0.26	76	0.23	54	2,23	86	1.20						
26	0.22			46	1.00	88	0.58						
34	0,59												
weighted average	0,28		0.16		0.72		1.04						

.

\$

٠

.

the Rung-Sat Special Zone might not detect the oncoming net, and therefore the catch in the Vung-Tau Region might be greater if 1-m plankton nets could be used.

Benthic Invertebrates. These are animals that live on or near the bottom or burrow into the mud, and include sponges, coelenterates, mollusks, polychaetes, echinoderms, and crustaceans. It is difficult to make direct comparisons of the kinds and numbers of benthos caught in the two study areas because a greater number and diversity of taxonomic groups occur at higher salinities. Nor can we necessarily infer greater productivity in comparing diversity of the number of different taxonomic groups in each region, simply because organisms such as sponges, corals, anemones, squids, oysters, starfish, and crabs are rare or absent in fresh water; these groups increase in numbers of individuals and species as the salinity regime increases (Fischer-Piette 1931).

In the Rung-Sat Special Zone, bottom trawls yielded 844 and 3012 invertebrates during the wet and dry seasons, respectively, while in the Vung-Tau Region there were 1552 and 1876 invertebrates during the wet and dry seasons, respectively (Table XII).

The large numbers of animals in the Rung-Sat are represented by only a few groups. Decapod shrimps formed the bulk of the collections from the Rung-Sat, representing 93 and 91 percent of the invertebrates during the wet and dry seasons, respectively; in the Vung-Tau collections they comprised 78 and 91 percent of the collections during the wet and dry seasons, respectively. In the Rung-Sat Special Zone during the wet season there was a predominance of crustaceans over a bottom salinity range of

Table XII.

Invertebrates and lower chordates collected with 10-ft otter trawl in mangrove region of SVN, 1972-73. Numbers in columns are number of specimens. The Rung-Sat is defoliated; Vung-Tau is nondefoliated (control) region. Wet-season collections were made in October and November 1972. Dry-season collections were made in January 1973. Dashes in columns indicate sample not collected.

RING-SAT	(defaliated)	 vet	50250D
V0140-9411	(derorrated)	 WCL	269900

VND stations	10	16	14	12	20	22	24	32	30	18	28	34	26	Total
Bottom salinity, ⁰ /00	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.5	23,9	26.5	26.7	
Porifera (sponges) Coelenterata	0	0	0	0	0	0	0	0	0	0	0	1 ²	0	1
Hydrozoa (hydroids)	0	0	0	0	0	0	1	0	0	0	0	0	0	1
Anthozoa (corals & anemones) Actinaria (anemones)	0	0	0	0	0	0	0	0	0	0	1	0	0	1
Madreporaria (stony corals) Annelida (worms)	0	0	0	0	0	2	0	0	0	0	0	1	0	3
Polyments (bristleworms)			Δ	٥	0	n	٥	٥	0	0	0	•	,	
l olyno == = 4	·		··¥			 ¥	· •	Ŭ	U	U	U	v	1	1
Syllidae	0	0	0	0	1	0	0	0	0	0	0	0	0	1
Nereidae	0	0	1	0	0	5	0	0	0	0	0	0	0	6
Goniadidae	0	0	0	0	1	0	0	0	0	0	0	0	0	1
Eunicidae	0	0	0	0	0	1	0	0	0	0	0	0	0	1
Orbiniidae	0	1	0	0	0	0	0	0	0	0	D	0	0	1
Unknown family Mollusca Gastropoda (snails)	0	4	0	0	0	0	0	0	0	0	0	0	0	4
mesogastropoda Cerithiidae (tide-flat shells)) 0	0	0	0	0	1	0	0	0	0	0	0	0	1
<u>Cerithidea</u> sp.	0	0	0	0	0	0	0	. 0	0	0	0	1	0	1

^a Fragments (considered as one individual)

ទទ

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations	10	16	14	12	20	22	24	32	30	13	28	34	26	Total
Bottom salinity, ⁰ /oo	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.5	23.9	26.5	26.7	
Calypta eid a e (cup & saucer sh <u>Calyptraea</u> sp.	e 11s) 0	0	0	0	0	0	0	0	0	0	0	0	1	1
<u>Crepidula</u> sp.	0	0	0	. 0	0	0	0	0	0	0	0	0	3	3
Naticidae (moon snails)	0	0	0	0	0	1	0	0	0	0	0	0	0	1
Natica sp.	0	0	0	0	1	0	0	0	0	0	0	0	0	1
cf. Bursa sp.	0	, 0	0	0	0	0	0	0	0	0	1	0	0	1
Neogastropoda Muricidae (rock shells)	0	0	0	0	0	0	6	0	0	0	0	0	0	6
Thais sp.	0	0	0	0	1	2	0	1	1	0	0	0	0	5
?Buccinidae (whelks)	0	0	0	0	0	0	0	0	1	0	0	0	0	1
?Mitridae (mitre shells) Pelecypoda (bivelves) Pteromorphia	0	0	0	0	0	0	0	0	1	0	0 ,	0	0	1
Arcidae (ark shells) <u>Arca</u> sp. Anomiidae (jingle shells)	. 0	0	0	0	0	0	0	0	1	0	. 0	0	0	1
Placuna placeota	0	0	0	0	0	0	0	0	0	0	0	1	0	1
Ostreidae (oysters)	0	0	0	0	0	0	0	0	0	0	0	· 1	0	1
"dead shells"	0	0	0	0	0	0	0	0	0	0	1	0	8	9
Cephalopoda (squid & cuttlefishes) Sepioidea (cuttlefishes) Arthropoda	0	0	0	0	0	0	0	0	0	0	0	1	0	1
Malacostraca (crabs, lobster, shr Stomatopoda (mantis shrimps) Squillidae	imps)													
<u>Squilla</u> sp. A	0	0	0	0	2	0	0	0	0	0	0	0	0	2

*

4

56

.

• .

-

•

۳

٠

.

RUNG-SAT (defoliated) -- wet season - (cont'd.)

•

.

.

ς.

VND stations	10	16	14	12	20	22	24	32	30	13	28	34	26	Total
Bottom salinity, ⁰ /00	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.5	23.9	26.5	26.7	
Mysidacea (opossum shrimps) Isopoda (peracarideans) Valvifera	0	0	0	0	3	0	0	0	2	4	0	0	0	9
Idoteidae	•		_	•	_	•		-			_		_	
Idotea sp.	Û	0	1	0	1	0	3	0	0	1	0	. 0	0	6
Amphipoda (Deach fleas)	~	~	•	~	•	•		•	~		~	•	~	_
Gammaricea Doceroda	U	U	U	v	0	U	U	U	U	T	Ų	U	U	1
Netentie (true shrimes)														
Panaeidet														
Pangaidaa														
Trachypenaeus sp.	0	Q	0	0	0	0	0	0	0	0	0	0	4	. 4
?Trachypenseus sp.	0	22	2	0	67	15	0	2	8	6	8	3	Ð	133
Unidentified genus	0	0	0	0	0	0	83	0	0	0	0	• 0	0	83
Sergestidae														
?Acetes sp.	0	0	2	1	11	1	10	0	51	372	0	0	1	449
Caridea (sand & snapping shri	mps)													
Alpheidae (snapping shrimps)	_	-	_	_	_		-	-	_	_	_	ĥ	`	
Alpheus sp.	1	0	1	0	0	0	0	0	Q	0	0	2~	1	5
Palaemonidae (prawns)		~	~		_	•	b	•	•		•	-		
Palaemontes sp.	10	0	0	U	0	0	11.	U	0	4	0	U	U	31
Macrobrachium sp.	4 ^b	10 ^b	0	0	7 ^b	9	4 ²	, O	0	0	0	0	0	34
Reptantia (lobster, crayfish,	crabs)													
Anomura (crabs)	•													
Diogenidae (hermit crabs)														
? <u>Clibanarius</u> sp. A	0	0	0	0	2	1	4	1	1	0	0	0	0	9
? <u>Clibanarius</u> sp. B	0	0	0	. 0	0	4	2	0	0	0	0	0	0	6

^bIncludes ovigerous specimens

.

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations	10	16	14	12	20	22	24	32	30	13	28	34	26	Total
Bottom salinity, ⁰ /00	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.	23.9	26.5	26.7	
Brachyura (true crabs) Portunidae (swimming crabs) ? <u>Portunus</u> sp.	0	0	0	0	0	0	1	0	0	U	0	0	0	1
<u>Charybdis</u> sp. Manthidao (mud orabs)	0	0	0	0	0	1	1	0	0	0	0	0	1	3
Unidentified genus Dorippidae	0	0	0	0	0	1	0	0	0	0	0	0	0	1
<u>Ethusa</u> sp. Leucosiidae	0	0	0	0	1	0	2	0	0	0	0	1	0	4
?Philyra sp.	0	0	0	0	0	0	0	0	2	0	0	0	0	2
Sipunculida (peanutworms) Echinodermata Asteroidea (sea stars) Phanerozonia	0	1	0	0	0	0	0	0	0	0	0	0	0	. 1
Luidiidae <u>Luidia</u> sp. Ophiuroidea (brittlestars)	0	0	0	0	0	0	0	0	0	0	1	0	0	1
Amphiuridae Ophiactidae	0	0	0	0	0	1	1	0	0	0	0	0	0	2
<u>Ophiactis</u> sp.	0	0	0	0	0	0	0	0	0	0	0	1	0	1
Total specimens	21	38	7	1	98	45	129	4	68	388	12	13	20	844

.

٠

4

58

-

.

.

· . •

~

.

RUNG-SAT (defoliated) -- dry sesson

F •

.

	VND stations	66	64	62	60	80	74	72	68	78	76	76	Total
	Bottom salinity, ⁰ /00	3.5	4.3	6.5	15.5	17,9	20.5	21.3	23,3	25 .9	26.1	27.3	
	Coelenterata		<u>,</u>		. 5				•				
	Hydrozoa (hydroids)	Ų	0	0	1	U U	Û	Û	U	0	I.a.	0	2
	Scyphozoa (medusae) Anthozoa (corals & anemones)	0	0	1	0	0	0	0	0	0	0	0	1
	Madreporaria (stony corals)	0	0	0	0	0	0	0	0	0	0	3	3
	Annelida (worms)	-	-	-	-	-	-	-	-	-	-	•	-
	Polychaeta (bristleworms) Eunicidae	0	0	0	0	0	0	0	0	0	0	4	4
თ	?Orbiniidae	0	0	0	0	0	1	0	0	0	0	0	1
Q	Sabellidae	0	0	240+	0	0	0	0	0	0	0	0	240+
	Unknown family Mollusca	0	0	3+	0	0	0	0	0	0	0	1	4 1
·	Gastropoda (snails)												
	Neogastropoda												
	Buccinidae (whelks)	0	0	1	0	0	0	0	0	0	0	0	1
	Marginellídae (marginellas)		_	_	_	_		_	_	_	_	_	
	<u>Hyalina</u> sp.	0	0	0	0	0	I	0	0	0	0	0	1
	Pelecypoda (bivalves)												
	Protobranchia												
	Muculidae (little nut shells)	,	0	0	0	٥	1	0	0	0	0	1	2
	<u>Mucutana</u> sp. Pteromorphia	v	v	v	0	v	T	0		v	•	1	2
	Mytilidae (masels)	0	0	1	0	0	0	0	0	0	0	1	2
	Heteroconchia	-	•	-	•	·	-	+	-	-	-	-	-
	Tellinidae (sunray clams)	0	0	0	0	0	1	0	0	0	0	0	1
	Aloididae	_							_				
	Corbula sp.	0	Ð	0	0	0	1	0	0	0	0	0	1

^a Fragments (considered as one individual)

•

• •

RUNG-SAT (defoliated) -- dry season - (cont'd.)

VND stations	66	64	62	60	80	74	72	68	78	70	76	Total
Bottom salinity, ⁰ /00	3.5	4.3	6.5	15.5	17.9	20,5	21.3	23.3	25.9	26.1	27.3	
Pholadidae (piddocks)			~		·		•		-	<u>^</u>		
Martesia sp.	U	0	U	0	U	Û	0	0	U	0	1	1
Unknown order Cephalopoda (squid & cuttlefishes) Sepioidea (cuttlefishes) Sepiidae	, ,	0	0	0	0	0	0	0	0	0	1	1
<u>Sepiella</u> sp. Arthropoda Crustacea	0	0	0	0	0	0	0	0	0	0	1	1
Malacostraca (crabs, lobster, shr	imps)											
Stomatopoda (mantis shrimps) Squillidae	0	0	0	0	0	0	0	0	0	1	0	1
<u>Squilla</u> ?quinquedentata	0	0	0	0	0	0	0	0	0	0	2	2
<u>Squilla</u> scorpio	· 0	0	0	0	1	0	0	0	0	0	0	1
Mysidacea (opossum shrimps) Isopoda (peracarideans) Flabellifera (parasitic) Sobastomidae	0	6	8	3	0	0	0	2	1	0	0	20
?Exosphaeroma sp.	0	5	0	0	0	0	0	0	0	2	0	7
? <u>Cilicaea</u> sp. Valvifera	Ø	· 0	0	0	0	0	0	0	0	0	1	1
Idoteidae <u>Cleantis</u> sp. Amphipoda (beach fleas) Cammaridea	0	0	0	0	0	1	0	2	0	0	0	3
?Talitridae (beach fleas)	0	0	4	0	0	0	0	0	0	0	0	4
Corophiidae	0	1	0	0	0	0	0	0	0	0	0	1

-

RUNG-SAT (defoliated) -- dry season - (cont'd.)

VND stations	66	64	62	60	80	74	72	68	78	70	76	Total
Bottom salinity, ⁰ /oo	3.5	4.3	6.5	15.5	17.9	20.5	21.3	23.3	25.9	26.1	27.3	

• ,

,

,

,

· . · ·	Decapoda Natantia (true shrimps) Penaeidea Penaeidae 	2	1	1	2	148	91	5	15	155	2	57	479
	Unidentified genus	-	0	0	0	0	0	0	0	0	4	0	4
	Sample												
	? <u>Acetes</u> sp.	1	312	233	353	100	16	346	261	57	298	111	2,088
	Pasiphaeidae	he)											
	Leptochela sp.	0	0	0	0	0	0	0	0	0	0	ვხ	3
	Alpheidae (snapping shrimps)			-			_	_					
61	<u>?Alpheus</u> sp.	0	1	3	0	0	0	0	0	0	0	1	5
	?Palaemon sp.	0	0	0	0	0	0	0	0	0	0	ക	6
		•	•	-	-	-	•	-	Ŭ	· ·	Ŭ	Ū	U
	?Leander sp.	0	0	0	0	2 ^b	0	0	0	0	0	0	2
	?Periclimenes sp.	0	0	0	o	0	0	0	0	2	0	3p	5
	Macrobrachium sp.	0	0	51 ^b	0	0	3 ^b	0	0	0	0	0	54
	Reptantia (lobster, crayfish, c	rabs)							-	_	_		
	Anomura												
	Porcellanidae (porcelain crab	s)	_	_	_								
	Unidentified genus	0	0	0	0	0	1	0	0	0	0	0	1
	Diogenidae (hermit crabs)	0	•	•	~	•	~	~		-	•	~	
	Paguristes sp.	Ŭ	0	0	0	U	U	0	U	2	U	U	2
	Unidentified genus sp. A	0	0	0	0	0	1	0	0	0	0	0	1
	Unidentified genus sp. B	0	0	0	0	0	1	0	0	0	0	0	1

^bIncludes ovigerous specimens

· · · ·

٠

RUNG-SAT (defoliated) -- dry season - (cont'd.)

VND stations	6 6	64	62	60	8 0	74	72	68	78	70	76	Total
Bottom salinity, ⁰ /00	3,5	4.3	6.5	15.5	17.9	20.5	21,3	23.3	25,9	26,1	27.3	
Brachyura (true crabs) Portunidae (swimming crabs) <u>Charybdis</u> ? <u>cruciata</u>	0	0	0	0	0	1	0	0	0	Ů	1	2
?Lissocarcinus sp.	0	0	0	0	0	1	0	0	0	0	0	1
Unidentified genus Vanthidae (mud.crahs)	0	0	0	0	1	0	0	0	0	0	0	1
?Leptodius sp.	0	0	1	0	0	0	0	0	0	0	0	1
? <u>Sphaerozius</u> sp. Dorippidae	0	0	36 ^b	0	0	0	0	0	0	0	0	36
Ethusa ?sexdentata	0	0	0	0	2	0	0	0	0	0	0	2
Ethusa sp.	0	0	0	0	0	4	0	0	1	0	0	5
? <u>Philyra</u> sp.	0	0	0	0	0	1	0	0	0	0	0	1
Unidentified family	0	0	0	0	0	0	0	1	0	0	0	1
Bryozoa (moss animals) Echinodermata Asteroidea (sea stars) Phanerozonia Inidiidae	0	0	0	0	0	0	0	1ª	0	0	0	1
Luidia sp.	0	0	0	0	0	0	0	0	1	0	0	1
Holothuroidea (sea cucumbers)	0	0	0	0	0	0	0	0	O	0	1	`1
Chaetognatha (arrow-worms)	0	0	0	0	0	0	0	1	0	0	1	2
Total specimens	3	326	583+	359	254	126	351	283	219	308	200	3,012+

٠

^aFragments (considered as one individual) ^bIncludes ovigerous specimens

٠

.

62

.•

VUNG-TAU (nondefoliated) -- wet season

	VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
	Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27,9	28.1	28.3	28,5	28,7	29.3	30.5	
	Porifera (sponges)	0	1 ^a	1ª	0	0	2 a	0	0	0	2	2	0	8
	Coelenterata Hydrozoa (hydroids)	0	1 ^e	o	0	0	0	0	0	0	0	4	0	5
	Scyphores (medusae)	· 0	0	0	0	0	0	0	0	0	0	1	0	1
	Anthozoa (corais & anemones) Pennatu lacea (sea pens) Ptorceididee	0	0	1	0	0	0	0	0	1	0	0	0	2
	<u>Pteroeides</u> sp.	0	0	1	13	1	0	0	0	8	0	0	8	31
	Actinaria (anemones)	0	0	0	0	0	0	0	0	0	2	1	2	5
	Nemertea (nemertean worms) Annelida (worms)	2	1 ^ª	0	0	Ó	0	0	19	0	1	0	0	5
თ	Polychaeta (bristleworms) Polynoidae	0	10	0	0	0	0	0	0	0	0	0	0	10
ىن	?Polynoidae	0	0	0	0	0	0	0	2	0	ł	0	0	3
	Phyllodocidae	1	0	0	0	0	0	0	2	0	0	0	0	3
	Hesionidae	0	3	0	0	0	0	0	1	0	0	4	0	8
	Syllidae	0	1	0	0	0	0	0	0	0	0	0	0	1
	Nereidae	0	2	0	0	0	0	0	1	0	3	1	0	7
	Goniadidae	3	0	0	0	0	0	0	1	0	0	o	0	4
	Onuphidae	0	0	Q	0	0	0	0	10	0	0	0	0	10

[®] Fragments (considered as one individual)

1

•

٠.

.

	VUNG	-TAU	(nonde	foliat	ed)	wet s	eason	- (cos	it'd.)				
VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27.9	28.1	28.3	28,5	28 7	29,3	30.5	
Eunicidae	2	2	0	0	0	<u>0</u>	0	0	0	0	0	1	5
Lumbrineridae	2	0	0	0	0	0	O	0	0	0	0	0	2
Orbiniidae	0	0	0	0	0	• 0	0	1	0	0	0	0	1
Opheliidae	0	- 1	0	0	0	0	0	1	0	0	0	0	2
Sternaspidae	1	0	0	0	0	3	0	0	0	0	0	0	4
Pectenariidae	0	. 0	0	0	0	0	0	0	0	1	0	0	1
Ampharetidae	0	1	0	0	0	0	0	0	0	0	0	0	1
Terebellidae	0	3	0	0	0	0	0	8	0	2	0	0	13
Sabellidae	0	17	0	0	0	0	0	0	0	0	2	0	19
Unknown family Mollusca	3	0	0	0	0	0	0	0	0	0	4	0	7
Gastropoda (snails) Mesogestropoda													
Calyptaeidae (cup & saucer s <u>Calyptraea</u> sp. Naticidae (moon snails)	hells) O	0	° O	0	0	0	0	0	0	0	0	1	1
Natica sp.	0	0	0	0	0	0	0	0	0	0	0	2	2
<u>Polynices</u> <u>didyma</u> Neogastropoda	0	0	0	0	0	0	0	0	0	0	0	1	1
Thais sp.	0	0	0	4	0	0	0	0	0	0	Û	0	4
Murex sp.	0	0	0	0	0	0	0	0	0	0	0	1	1

.

.

. 8

,

۰.

,

\$

.

*

.

	VUNG	-TAU	(nonde	foliat	ed)	wet s	eason	- (con	t'd.)				
VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27.9	28.1	28,3	28.5	28.7	29.3	30.5	
Mitridae (mitre shells)	6	0	0	0	0	0	0	0	0	4	Ũ	0	10
?Mitridae	0	0	0	0	0	0	0	3	0	0	0	0	3
Volutidae (volutes)	0	0	0	1	0	0	0	0	0	0	0	0	1
Nudibranchia (sea slugs)	0	0	1	0	0	• 0	0	0	0	0	0	0	1
Unknown order Scaphopoda (tooth shells)	0	0	0	0	0	6	0	0	0	0	0	0	6
Dentaliidae Pelecypoda (bivalves) Protobranchia	0	0	0	0	0	0	0	0	0	0	0	1	1
Nuculidae (little nut shells) <u>Nuculana</u> sp. Pteromorphia	0	0	0	0	0	3	0	2	0	6	0	0	11
Arcidae (ark shells)	0	0	0	0	0	1	0	0	0	0	0	0	1
<u>Arca</u> sp. Mytilidae (muasels)	0	0	0	1	0	0	0	0	0	2	0	0	3
Brachydontes sp. Limidae (file shells)	7	0	0	0	0	0	0	3	0	4	0	0	14
Lima sp. Anomiidae (jingle shells)	0	0	0	0	0	0	0	1	0	0	0	0	1
Anomia sp. Heteroconchia	0	0	0	0	0	0	0	0	0	0	0	3	3
? <u>Cumingia</u> sp.	0	0	0	1	0	0	0	6	0	0	0	0	7
Solenid ae (razor c lams)	2	0	0	0	0	0	0	10	0	0	0	0	12
Pholadidae (piddocks)	0	0	1	0	0	0	0	0	0	0	0	0	1.
Unknown order	0	0	0	0	0	0	0	?+	0	12	0	3	15+

.

VUNG-TAU (nondefoliated) -- wet season - (cont'd.)

VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27,9	28.1	28.3	28,5	28 7	29,3	30.5	
"dead shells" Corbologoda (squid & cuttlefishes)	0	0	0	3	0	0	0	0	0	D	0	0	3
Sepioidea (cuttlefishes)	0	Ō	Ð	0	0	1	0	0	0	0	0	0	1
<u>Sepiells</u> sp. Arthropoda Merostomata Xiphosura (horseshoe crabs)	0	0	0	1	0	0	0	0	0	0	0	0	1
<u>Carcinocorpius</u> <u>rotundicauda</u> Crustacea Cirrepedia (barnacles)	1	0	0	0	0	0	1	0	0	0	0	0	2
Balanidae <u>Balanus</u> sp. Malacostraca (crabs, lobster, shr Stomatopoda (mantis shrimps)	0 imps)	0	0	1	0	0	1	0	0	. 1	0	0	3
Squilla fascieta	0	0	0	0	0	0	0	0	0	0	0	1	1
<u>Squilla</u> sp. A	0	0	0	0	0	0	0	0	0	0	1	Û	1
<u>Squilla</u> sp. B	0	0	0	0	0	0	0	0	1	0	0	0	1
<u>Squilla</u> sp. C	0	0	. 0	0	0	0	3	0	0	0	0	0	3
Mysidacea (opossum shrimps) Isopoda (peracarideans) Valvifera Idoteidae	20	5	0	31	0	0	4	19	0	32	15	1	127
<u>Idotea</u> sp. Epicaridea (parasitic) Bopyridae	2	0	0	0	0	0	0	0	0	0	0	0	2
?Bopyrina sp.	0	0	0	0	0	0	0	1	0	0	0	0	1
Orbione sp.	0	0	0	1	0	0	0	0	0	0	0	0	1

3

7

λ.

•

•

.

-

VUNG-TAU (nondefoliated) -- wet season - (cont'd.)

VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27.9	28.1	28.3	28.5	28.7	29,3	30.5	
Amphipoda (beach fleas)													
Gammaridea	0	2	0	0	0	0	1	0	0	1	0	0	4
Decapoda													
Natantia (true shrimps)													
Penaeidea													
renae10ae	110	•	10	0.1		1.05	A E			•	-		
racnypenaeus sp.	119	Z	12	21	15	105	35	111	17	8	3	10	458
Acctos and	1.05	Ω	0	ń	0	160	14	6	1	1.1		•	000
Curidea (cand & examples shri		U	U	v	Ŷ	102	14	Ŷ	1	41	T	2	332
Alphaidaa (enapping shrimpa)	անցչ												
741 nhave en	6b	gb	70	0	0	0	1	1	4	0	0	0	27
Palaemonidae	Ŷ	Ŭ	,	ĩ	Ť	•	-		-	v	v	v	21
Palaemonetes sp.	0	0	0	0	0	0	0	0	۷p	' O	0	0	4
	•	•	•	•	-	•	•	-	-	•	•	v	-
?Palæemonetes sp.	13 ^b	0	0	27 ^b	0	0	3 ^b	43 ^b	0	3p	0	0	89
Macrobrachium sp.	0	1	0	0	0	0	0	0	0	0	0	0	1
Unidentified genus	0	0	0	0	0	0	0	0	ıb	0	0	0	1
Reptantia (lobster, crayfish,	crabs)												
Anomura													
Porcellanidae (porcelain crai	bs)	•	.b	- oh	~	•	•	-	h	-	-	- b	
Porcellana sp.	Ų	U	L	135	U	U	U	U	13*	0	U	70	34
Unidentified genus	0	3	0	0	0	0	Ð	0	0	0	0	0	3
Diogenidae (hermit crabs)												ъ	
? <u>Clibanarius</u> sp. C	0	0	0	. 0	0	0	0	0	0	0	0	20	2
Brachyura (true crabs)													
Fortunidae (swimming crabs)	^	•	•	b	~		-				-		
rortunus sp.	U	U	U	3-	U	U	U	0	1	Q	0	3	7

^b Includes ovigerous specimens

67

• •

ς.

VUNG-TAU (nondefoliated) -- wet season - (cont'd.)

VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰	/00 26.7	27.1	27.3	27.5	27.7	27.9	28.1	28.3	28.5	28.7	29,3	30,5	
Charybdis ?natator		0	0	0	0	0	10 ^{°t}	0 10	0	0	3	0	23
<u>Charybdis</u> sp.	C) 0	0	17 ^b	° 1	0	0	0	0	0	0	0	18
Scylla serrata	() O	0	1	0	0	0	0	0	0	0	0	1
Podophthalamus sp.	e) O	0	1	0	0	0	0	0	0	0	0	1
Xanthidae (mud crabs) ? <u>Myomenippe</u> sp.	c	0	0	1	o	0	0	0	0	0	0	0	1
Unidentified genus	c) 1	1	0	0	0	0	0	0	0	0	0	2
Majidae (spider crabs) Unidentified genus	c) 0	0	1	0	0	1	0	0	0	0	0	2
Dorippidae <u>Ethusa</u> sp.	1	. 2	0	4	. 2	3	1	12	0	7	1	5 ^b	38
Leucosiidae ? <u>Philyra</u> sp.	1	. 0	0	0	0	0	2	1	0	0	0	0	4
Unidentified family	c) o	0	2	0	1	0	10 ^t) O	0	0	0	13
Sipunculida (peanutworms) Echinodermata	c	0	0	0	0	0	0	0	0	0	0	2	2
Ophiuroidea (drittlestars) Ophiurae Amphiuridae	1	. ⁸ 1	0	0	0	2	0	4	0	1	0	0	9
Ophiotrichidae Echinoidea (sea urchins) Centrechinoidea	C	8	2	0	0	0	0	0	0	4	0	0	14
Temnopleuridae <u>Salmacis</u> dussumieri	C	0	0	1	0	0	0	0	0	. 0	6	0	7

.

-

^aFragments (considered one individual) ^bIncludes ovigerous specimens

*

.

68

.
	VC	ING-TAU	(non	defoli	ated)	wet	seaso	on - (c	ont'd.)			
VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, 0/00	26.7	27.1	27.3	27.5	27.7	27.9	28.1	28.3	28.5	28.7	29.3	30.5	
Holothuroidea (sea cucumbers)	0	23	1	2 ^a	2	0	0	0	0	23	0.	0	51
Ascidiacea (sea squirts)	0	2	0	0	0	0	0	0	0	2	0	0	4
Total specimens	298	99	29	151	21	289	77	271+	51	161	49	56	1,552+

.

^aFragments (considered as one individual)

2

VUNG-TAU (nondefoliated) -- dry season

VND stations	100	84	104	82	86	90	88	102	92	98	94	96	Total
Bottom salinity, 0/00	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32.1	323	33.3	33.5	
Porifera (sponges) Comlenterata	0	0	0	4 a	O	0	2	0	0	э	-	0	6
Hydrozoa (hydroids)	0	0	1	4	1	1+	1+	• •	0	0	-	0	8+
Hydroida (hydrozoans) Sertulariidae	0	0	0	0	0			0	0	0	-	2	41
Scyphozos (medusae) Anthozos (corals & anemones) Telestacea Telestidae	0	1	0	0	0	1	0	1	0	0	-	1	4
<u>Telesto</u> sp. Gorgonacea (horny corals) Melitodidae	1	0	0	0	1	0	1	0	0	3	-	3	9
Melitodes sp.	0	0	0	0	0	0	0	0	0	0	-	1	1
Pennatulacea (sea pens) Pteroceididae	0	0	0	0	0	0	0	1 ²	0	0	-	0	1
Pteroeides sp.	2	0	0	1	0	0	0	0	0	0	-	0	· 3
Actinaria (anemones) Annelida (worms)	1	0	0	0	0	0	0	0	0	0	-	1	2
Polychaeta (bristleworms) ?Polynoidae	0	0	0	0	0	18	19	0	0	0	-	. 0	20
Hesionidae	0	0	0	1	0	0	1	0	0	0	-	0	2
Nereidae	0	0	0	8	0	0	5	0	0	0	-	0	13
Eunicidae	0	4	0	2	1	0	2	0	0	0	-	0	9
Terebellidae	0	0	• 0	0	0	0	17	0	Û	0	-	0	17
Unknown family	0	0	0	0	0	0	4.	0	0	la	-	Û	5

.

٠

²Fragments (considered as one individual)

a •

.

• • • •

		VUNG-	TAU ((nondei	oliate	ed)	dry se	ason -	(cont	'd.)				
	VND stations	100	84	104	82	8 6	90	88	102	92	98	94	96	Total
	Bottom salinity, ⁰ /00	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32.1	32.3	33,3	33,5	
	Mollusca													
	Gastropoda (snails)													
	Trochidae (top shells)													
	Euchelus sp.	0	0	0	0	0	0	1	0	0	0	-	0	1
	Mesogastropoda													
	Turritellidae Turitella bacillum	n	٥	0	0	Δ	0	0	•	0	,		•	1
	Idriteria Dacirium	Ý	U	v	v	v	0	U	v	v	1	-	U	1
	Lamellaridae	0	0	0	0	0	0	1	0	0	0	-	0	1
	Neogastropoda													
4	Muricidae (rock shells) Murex ternispina	0	0	0	0	0	0	0	0	1	0	-	0	1
	Columbellidae		-	-	-	-	-	•	•	-	•		•	-
	Anechis sp.	0	0	0	0	0	0	1	0	0	0	-	0	1
	Fasciolariidae (tulip sheils) Latirus so	. 0	0	0	0	0	0	0	0	0	0	_	1	,
	Laciida op.	0	Ŷ	Ť	v	Ŭ	v	v	Ý	v	v	-	I	±
	Nudibranchia (sea slugs)	0	0	0	0	ò	0	1	0	0	0		0	1
	Pelecypoda (bivalves)													
	Nuculidae (little nut shells)	0	0	0	1	0	0	0	0	o	0	-	0	1
														_
	<u>Nucula</u> sp.	0	0	0	0	0	0	1	1	0	0	-	0	2
	Arcidae (ark shells)	0	0	0	1	0	0	0	0	0	0	-	0	3
		-	-	-	-	-		-	-	-	•		·	-
	<u>Anadara</u> sp.	0	0	0	2	0	0	1	0	0	0	-	0	3
	Mytilidae (mussels)	0	0	0	0	0	0	1	0	0	0	-	0	1
	Protividas (seallest)	0	~	0	,	0	^	0	0	~	0		•	1
	Limidae (file shells)	0	v	U	Ŧ	9	0	U	0	v	v	-	U	1
	Lima sp.	Ú.	· 0	0	0	0	0	1	0	0	0	-	0	1

.

· · · ·

VUNG-TAU (nondefoliated) -- dry season - (cont'd.)

VND stations	100	84	104	82	86	90	88	102	92	98	94	96	Total
Bottom salinity, ⁰ /oo	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32.1	32,3	33.3	33.5	
Anomiidae (jingle shells)													
<u>Enigmonia</u> sp.	0	1	0	0	0	0	0	0	0	0	-	0	1
Ostreidae (oysters)	0	0	0	0	0	0	1	0	0	0	-	0	1
<u>Crassostrea</u> sp. Heteroconchia	0	0	0	0	0	0	1	0	0	0	-	0	1
Veneridae (clams)					•								
Pitar sp.	0	1	. 0	0	0	0	0	. 0	0	0	-	0	1
Tellinidae (sunray clams) Solenidar (razor clams)	0	0	0	0	0	0	1	0	0	0	-	0	1
<u>Solen</u> sp.	0	1	0	0	0	0	0	0	0	0	-	0	1
Saxicavidae (boring clams)	0	0	0	0	0	0	1	0	0	0	-	0	1
Cephalopoda (squid & cuttlerishes) Teuthoidea (squid) Sepioidea (cuttlefishes) Sepiidae	0	0	0	1	0	0	0	0	0	0	•	0	1
<u>Sepia</u> sp.	0	0	0	1	0	0	0	0	0	0	-	0	1
<u>Sepiella</u> sp. Arthropoda	0	0	0	0	0	0	0	0	0	0	-	2	2
Crustacea	_		_	_		_	_	_	_				
Ostracoda (seed shrimps) Cirrepedia (barnacles) Balanidae	0	0	0	0	0	0	0	0	0	0	-	2	2
<u>Balanus</u> sp. Malacostraca (crabs, lobster, shr	0 imps)	1	1	0	0	2	0	0	0	0	-	0	4
Stomatopoda (mantis shrimps)	- •												
Squillidae	0	0	8	0	7	0	0	0	0	0	-	0	15
<u>Squilla</u> ? <u>quinquedentata</u>	0	0	0	1	0	4	0	0	0	0	-	0	5

.

VUNG-TAU (nondefoliated) -- dry season - (cont'd.)

.

VND stations	100	84	104	82	86	90	88	102	92	9 8	94	96	Total
Bottom salinity, ⁰ /00	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32.1	32.3	33,3	33.5	
<u>Squilla scorpio</u>	0	0	1	0	0	0	0	0	0	Û	-	0	1
<u>Squilla</u> sp. A	0	0	0	0	0	0	0	0	0	0	-	1	1
Mysidacea (opossum shrimps) Isopoda (peracarideans) Flabellifera (parasític) Sobseromidae	0	8	6	20	57	30	744	0	1	0	-	15	881
? <u>Cilicaea</u> sp. Epicaridea (parasitic)	0	0	0	1	0	0	0	0	0	0	-	0	1
Epirenaeon sp.	0	0	0	0	0	0	1	0	0	0	-	0	1
Probopyrus sp. Amphipoda (beach fleas)	0	1	0	0	0	0	0	0	0	0	-	0	1
Gammaridea ?Talitridae (beach fleas)	0	0	0	3	0	0	31	0	0	0	*	0	34
Corophiidae Decapoda Natantia (true shrimps) Penaeidea	0	6	0	13	14	0	1	0	0	0	-	0	34
Penaeidae ?Trachypenaeus sp.	4	20	12	2	13	0	13	8	2	3	-	10	87
Sergestidae ? <u>Acetes</u> sp. Caridea (sand & snapping shrim	0 aps)	1	10	0	0	1	23	l	2	2	-	384	424
Hippolytidae ? <u>Lysmata</u> sp.	0	0	0	0	0	0	1	0	0	0	-	2	3
Alpheidae (snapping shrimps) ? <u>Alpheus</u> sp.	C) 1	Û	2	0	0	4 ^t	° 0	0	0	-	0	7

^bIncludes ovigerous specimens

VUNG-TAU (nondefoliated) -- dry season - (cont'd.)

VND stations	100	84	104	82	86	90	88	102	9 2	98	9 4	96	Total
Bottom salinity, ⁰ /00	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32,1	32. 3	33.3	33.5	
Palaemonidae				•									
? <u>Leander</u> sp.	0	40 ^b	6 b	0	1	0	1	0	0	0	-	дþ	57
Unidentified genus ?Gnathophyllidae	lp	0	0	0	0	0	0	0	0	0	-	0	1
Unidentified genus	0	0	0	0	0	0	0	0	0	0	-	1	1
Reptantia (lobster, crayfish, c:	rabs)												
Ancoura													
Porcellanidae (porcelain crab <u>Porcellana</u> sp.	s) 5 ^b	0	0	2	0	0	4	0	0	Û	-	1	12
?Porcellana sp.	0	0	0	0	24	2	0	0	0	0	-	0	26
Megalobrachium sp. Brachyura (true crabs) Portunidae (swimming crabs)	0	0	0	0	0	0	1 ^b	0	0	0	-	0	1
Portunus sp.	0	0	5	0	1	0	1	6	0	0	-	0	13
<u>Charybdis</u> ?cruciata	0	2	0	0	· 7	0	3	3 ^b	7 ^{°C}	, o	-	8 ^b	30
?Lissocarcinus sp.	0	0	0	1	1	0	0	0	0	0	-	0	2
Scylla serrata	o	0	0	0	0	0	0	0	0	0	-	1	1
Unidentified genus	1	0	0	0	0	0	0	0	0	0	-	0	1
Micropanope sp.	0	0	0	0	0	1	47 ^b	0	0	0	-	0	48
Unidentified genus Maiidae (spider crabs)	0	0	0	0	0	0	0	0	0	0	-	1	1
<u>Pugettia</u> sp.	0	0	I	1	0	0	0	0	0	0	-	0	2
? <u>Hyastenus</u> sp.	0	0	0	0	0	0	0	0	0	0	-	1	1

^b Includes ovigerous specimens

74

	Vt	NG-TAU	(nor	ndefoli	iated)	dry	Seaso	n - (c	ont'd.	>			
VND stations	100	84	104	82	86	90	88	102	92	9 E	. 94	96	Total
Bottom salinity, ^C	/00 30.5	30.5	30.7	30.7	30.7	30.9	31,1	31.7	32.1	32,3	33,3	33.5	
? <u>Maja</u> sp.	0	0	0	0	0	0	0	0	Ō	0	-	0	1
Dorippidae Dorippe ?granulata	1	0	0	0	0	0	0	0	0	0	-	0	1
Dorippe sp.	0	0	0	0	0	0	0	0	0	0	-	1	1
Ethusa sp.	0	2	0	5	0	0	1	2	1	0	-	0	11
Bryozoa (moss animals) Echinodermata Asteroidea (sea stars) Phanerozonia	0	0	0	0	0	0	1ª	12	18	1 0 	-	0	3
Luidiidae Luidia sp. Ophiuroidea (brittle stars)	0	0	0	0	0	0	3	0	1	. 0	-	0	4
Ophiotrichidae Echinoidea (sea urchins) Centrechinoidea	0	3	0	3	0	0	14	0	0	0	-	2	22
Temnopleuridae <u>Salmacis dussumieri</u> Holothuroidea (sea urchins) Apoda	0	0	0	0	0	0	0	0	0	0	-	1	1
Caudinidae ? <u>Aphelodactyla</u> sp.	0	0	0	0	0	0	1	0	0	0	-	0	1
Chaetognatha (arrow-worms)	0	0	0	0	0	0	0	0	1	0	-	0	1
Chordata Ascidiacea (sea squirts)	0	1	0	0	0	0	2	0	1	0	-	0	4
Total specimens	16	9 3	51	81	128	44+	96 0+	24	17	10	-	452	1,876+

.

.

⁸Fragments (considered as one individual)

.

75

٠

.

13 to $18^{\circ}/\circ\circ$, mainly decaped shrimp, whereas their number greatly increased during the dry season, over a salinity range of about 4 to $27^{\circ}/\circ\circ$. Crustaceans occurred at every station in the Vung-Tau Region. There is a greater diversity, as reflected in diversity indices of taxa, of animals in the Vung-Tau Region both during the wet season (2.21) and dry season (1.87) than in the wet season (1.69) and dry season (0.92) in the Rung-Sat Special Zone. Thus we may infer the Rung-Sat Special Zone represents an impoverished, stressed ecosystem, in contrast to the relatively rich invertebrate fauna in the Vung-Tau Region.

Because of the strong tidal currents in the Rung-Sat Special Zone as compared to the Vung-Tau Region, we might expect benthonic animals to experience more difficulty in finding an adequate habitat in which to hide or on which to hold in the Rung-Sat. However, because of the large amount of leaf, bark, and wood debris found along the estuarine bottom of the Rung-Sat Special Zone, we initially believed that large numbers of organisms might be found using this newly-made habitat for shelter. On the contrary, samples with greater amounts of debris usually had fewer organisms than did samples with less debris. Unfortunately, we were unable to quantify these data because of the difficulty in obtaining statistics on the area over which the trawl fished in both study areas. It is surmised that the large amounts of leaf, bark, and wood on the bottom of the Rung-Sat Special Zone represent a region inhospitable to many kinds of organisms because of the bacterial decomposition. This process would remove oxygen and could add toxic hydrogen sulfide to the interface between the bottom muds and the decomposing debris. Although we could not readily detect

the smell of hydrogen sulfide in samples of debris, there was usually a distinct smell of decomposition, even though two years had passed since the defoliants were sprayed. A letter from Mr. Huynh-trung-Hat, Fishery Specialist for Vung-Tau City, to the Director of Fisheries, Saigon, stated:

When the Allied Forces were spraying defoliants [1962-70], practically all the vegetation (approximately 95 percent) in Ban and Duoc mangroves were killed. As a result, the dead foliage caused the water to stink and the fauna (especially shrimp) to move to other areas.

Because we were probably unable to sample the oxygen content of the interface immediately between the bottom muds and the debris overlying them, it is not possible to show if low oxygen content might be a factor limiting the distribution and abundance of benthos in the Rung-Sat Special Zone.

Fishes. Juvenile and adult fishes were collected with a 10-ft bottom trawl (Table XIII). In the Rung-Sat Special Zone, 104 specimens of 29 species were taken in the wet season and 215 specimens of 17 species were caught in the dry season. Vung-Tau collections yielded 240 specimens of 34 species during the wet season, and 250 specimens comprising 22 species in the dry season. Diversity indices of fish families (Table XIV) show that the Rung-Sat Special Zone reflects the richest catches during the wet season, and the most impoverished fauna during the dry season, while the fish fauna of the Vung-Tau Region was intermediate, and remained similar during both seasons. Presumably the wide seasonal variation in diversity in the Rung-Sat Special Zone is due to seasonal salinity variation. If species diversity indices are examined (Table XV), the richest fish fauna occurred during the wet season in both study areas, with the Rung-Sat Special Zone having the

Table XIII.

Fishes collected with 10-ft otter trawl in mangrove region of SVN, 1972-73. Numbers in columns are numbers of specimens. The Rung-Sat is defoliated; Vung-Tau is nondefoliated (control) region. Wet season collections were made in October and November 1972. Dry season collections were made in January 1973. Dashes in columns indicate sample not collected.

			RUNG	G-SAT ((defoli	ated)	wet	seaso	n					
VND stations	10	16	14	12	20	22	24	32	30	18	28	34	26	Total
Bottom salinity, ⁰ /00	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.5	23.9	26.5	26.7	
<u>Físhes</u> Dasyatidae (stingrays)														
Dasyatis ?kuhlii	0	0	0	0	0	0	1	0	0	0	0	0	0	1
Urolophoides sp.	0	0	0	0	2	1	0	0	0	0	0	0	0	3
Engraulidae (anchovies) <u>Coilia</u> grayi	0	0	0	0	1	0	0	0	Û	4	0	0	0	5
Synodontidae (lizardfishes) <u>Harpodon</u> nehereus	0	0	0	0	0	0	1	0	0	0	0	0	0	1
Ariidae (sea catfishes) Arius macronotacanthus	4	8	0	0	0	0	1	0	0	0	0	0	0	13
Plotosidae (catfish eels) <u>Plotosus</u> <u>canius</u>	0	0	0	0	0	0	0	O	o	0	1	0	0	1
Polynemidae (threadfins) Polynemus longipectoralis	0	1	0	0	0	0	1	0	0	0	0	0	0	2
Apogonidae (cardinalfishes) Apogon quadrifasciatus	0	o	0	0	0	o	0	0	0	0	0	0	2	2
Lutjanidae (snappers) Lutjanus johni	0	0	0	0	0	0	1	1	0	0	0	0	0	2
Lutjanus ?johni	0	0	0	0	1	0	0	0	0	0	0	0	0	1

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations	10	16	14	12	20	22	24	32	30	18	28	34	26	Total
Bottom salinity, ⁰ /00	0.2	0.2	2.7	3.3	13.3	14.1	14.5	17.3	17.9	18.5	23.9	26.5	26.7	
<u>Fishes</u> Leiognathidae (ponyfishes) <u>Leiognathus berbis</u>	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Leiognathus insidiator	0	0	0	o	0	0	0	0	0	Ð	1	0	0	1
Gerreidae (mojarras) <u>Gerres</u> <u>abbreviatus</u>	0	0	0	0	0	0	0	0	0	0	1	0	0	1
Pomadasyidae (grunts) Pomadasys hasta	0	0	0	0	0	0	0	1	0	0	1	0	0	2
Sciaenidae (croakers) ? <u>Nibea acuta</u>	0	1	1	1	0	Û	1	0	0	4	0	o	0	8
Pseudosciaena crocea	0	0	0	0	0	0	1	0	0	0	0	0	0	1
?Wak sina	0	0	0	0	10	13	7	0	1	0	4	0	0	35
Eleotridae (sleepers) Prionobutis koilomatodon	[.] 0	0	0	0	0	0	1	0	0	0	0	0	0	1
Gobiidae (gobies) <u>Acentrogobius</u> <u>caninus</u>	0	0	0	0	0	0	o	1	0	0	0	0	1	2
Acentrogobius sp.	0	1	0	0	0	0	0	0	0	0	0	0	0	1
Apocryptodon cf. madurensis	0	1	0	0	0	0	0	0	0	0	1	0	0	2
Ctenogobius sp.	0	0	0	0	0	1	0	0	0	0	0	0	0	1
Glossogobius giurus	0	0	0	0	0	1	0	0	1	0	0	0	0	2
Trypauchen vagina	0	1	0	0	1	0	0	0	0	0	0	0	Q	2
Unidentified genus	0	0	0	0	0	1	0	õ	0	0	0	0	0	1

.

RUNG-SAT (defoliated) -- wet season - (cont'd.)

VND stations	10	16	14	12	20	22	24	32	30	18	28	34	26	Total
Bottom salinity, 0/00	0.2	0.2	2.7	3.3	13.3	14.1	14,5	17.3	17.9	18 5	23.9	26.5	26.7	
Cynoglossidae (tonguefish)	0	0	0		2	•		•					•	
Cynoglossus farei	0	U	U	U	3	U	U	0	U	0	0	0	0	3
Cynoglossus lingua	0	0	0	0	4	0	1	0	0	0	0	0	0	5
Soleidae (soles) <u>Zebrias</u> sp.	0	2	0	0	0	0	0	O	0	0	0	0	0	2
Batrachoididae (toadfishes) Batrachus grunniens	1	0	0	. 0	0	1	0	0	0	0	0	0	0	2
Total specimens	5	15	1	1	22	18	16	3	2	8	9	0	4	104

λ.

•

.

.

- ---

RUNG-SAT (defoliated) -- dry season

-

	VND stations	66	64	62	60	80	74	72	68	78	70	76	Total
	Bottom salinity, ⁰ /00	3.5	4.3	6.5	15.5	17.9	20.5	21.3	23.3	25.9	26.1	27.3	
	<u>Fishes</u> Clupeidae (h erring s) ? <u>Harengula</u> ovalis	0	0	0	0	0	0	0	1	0	0	o	1
	? <u>Sardinella</u> sp.	0	0	0	1	0	0	0	. 0	1	0	1	3
	Engraulidae (anchovies) <u>Coilia grayi</u>	0	1	2	0	3	7	7	3	3	1	1	28
	Ariidae (sea catfishes) Arius macronotacanthus	1	0	9	0	0	1	0	0	6	0	0	17
	Polynemidae (threadfins) Polynemus longipectoralis	0	1	1	0	0	0	0	0	0	0	0	2
18	Gerreidae (mojarras) <u>Gerres filamentosus</u>	0	0	0	0	0	0	0	0	0	0	1	1
	Pomadasyidae (grunts) <u>Pomadasys</u> hasta	0	0	0	0	0	1	0	0	0	0	0	1
	Sciaenidae (croakers) <u>Pseudosciaena</u> crocea	0	0	0	O	0	0	0	0	1	1	0	. 2
	<u>Sciaena</u> <u>russelli</u>	0	0	0	0	0	1	0	0	0	0	1	2
	? <u>Wak</u> <u>sina</u>	0	0	0	0	6	0	2	3	125	0	8	` 144
	Gobiidae (gobies) Parachaeturichthys polynema	0	0	0	0	1	0	0	0	0	0	0	1
	Trypauchen vagina	0	0	0	0	0	1	0	0	0	0	0	1
	Unidentified genus	0	0	0	0	0	0	0	0	1	0	0	1

	RUNG	-SAT (d	lefolia	ated) ·	dry	season	1 - (cc	ont'd.)				
VND stations	66	64	62	60	80	74	72	68	78	70	76	Total
Bottom salinity, ⁰ /00	3.5	4.3	6.5	15.5	17.9	20.5	21.3	23.3	25.9	26.1	27.3	
<u>Fishes</u> Scorpaenidae (scorpionfishes) <u>Leptosynaceia</u> asteroblepa	0	0	0	0	o	1	1	0	0	J	0	2
Cynoglossidae (tonguefish) Cynoglossus brachycephalus	0	Ū	0	0	0	0	0	0	0	0	1	1
Cynoglossus lingua	0	0	0	0	1	6	0	0	0	0	0	7
Cynoglossus puncticeps	0	0	0	0	1	0	0	0	0	0	0	1
Total specimens	1	2	12	1	12	18	10	7	137	2	13	215

· .

VUNG-TAU (nondefoliated) -- wet season

VND stations	42	44	40	46	48	56	58	50	38	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27.9	28,1	28.3	28.5	28.7	29.3	30.5	
<u>Fishes</u> Dasyatidae (stingrays) <u>Dasyatis</u> ?kuhlii	0	0	0	1	0	0	1	0	0	0	0	0	2
Centropomídae (snooks) <u>Ambassis</u> <u>dayi</u>	0	. 0	0	0	o	2	0	0	0	0	0	0	2
Apogonidae (cardinalfishes) Apogon <u>quadrifasciatus</u>	0	0	1	1	5	3	1	0	8	2	0	3	24
Sillaginidae (whitings) Sillago ?sihama	10	0	0	0	0	0	0	o	0	0	0	0	10
<u>Sillago</u> sp.	0	l	0	0	0	0	0	0	0	0	0	0	1
Lutjanidae (snappers) Lutjanus johni	1	1	1	0	3	0	0	0	0	0	0	0	6
Leiognathidae (ponyfishes) Leiognathus berbis	3	0	0	0	0	0	0	0	0	0	0	0	3
Leiognathus insidiator	8	0	0	0	0	1	0	0	0	0	0	0	9
Leiognathus ?insidiator	0	0	2	0	0	0	0	0	Ð	0	0	0	2
Pomađasyidae (grunts) Pomađasys hasta	13	. 2	0	1	1	12	0	4	0	3	0	1	37
Sciaenidae (croakers) <u>Sciaena russelli</u>	2	0	٥	0	0	1	0	8	0	0	0	0	11
? <u>Wak sina</u>	0	0	0	4	1	0	3	0	0	1	3	0	12

.

.

.

VUNG-TAU (nondefoliated) -- wet season - (cont'd.)

VND stations	42	44	40	46	48	56	58	50	3 8	52	54	36	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27.5	27.7	27.9	28.1	28.3	28.5	28.7	29.3	30. 5	
Fishes Callionumidae (daragenete)													
Calliurichthys fluviatilis	0	0	0	0	O	0	0	0	0	0	0	2	2
Calliurichthys japonicus	3	0	0	0	0	7	0	4	1	0	0	0	15
Eleotridae (sleepers) Pricnobutis koilomatodon	2	1	· 0	0	0	3	0	1	6	. 4	0	2	19
Gobiidae (gobies) <u>Acentrogobius</u> caninus	9	0	0	1	0	0	-3	0	0	0	0	0	4
Acentrogobius triangularis	0	0	0	0	0	6	0	0	0	0	0	0	6
Bathygobius fuscus	0	0	Ö	. 0	1	0	0	0	0	0	0	0	1
Bathygobius ?fuscus	0	0	0	0	0	0	0	1	0	0	0	0	1
<u>Bethygobius</u> fuscus	0	0	0	1	0	0	0	0	0	0	0	0	1
Cryptocentrus russus	0	0	0	0	0	0	0	2	0	0	0	0	2
<u>Glossogobius</u> giurus	12	0	0	0	0	20	0	0	0	0	0	1	33
Trypauchen vagina	0	0	0	0	1	0	0	0	0	0	0	0	1
Unidentified genus	O	O	0	0	0	1	0	0	0	0	0	0	1
Scorpaenidae (scorpionfishes) <u>Minous monodectylus</u>	0	0	0	1	0	0	0	0	0	0	0	1	2

ŧ

84

• .

	VUNG -	-TAU (nondef	oliate	d)	wet se	ason -	(cont	'd.)				
VND stations	42	44	40	46	48	56	58	50	38	52	54	3 6	Total
Bottom salinity, ⁰ /00	26.7	27.1	27.3	27,5	27.7	27 .9	28.1	28.3	28.5	28,7	29.3	30,5	
<u>Fishes</u> Platycephalidae (flatheads) Grammoplites scaber	0	0	0	8	0	0	ŧ	0	0	0	0	n	a
Rethides (loft-head floundone)	-	-	·	•	-	•	*	Ū	Ū	Ū	Ū	v	,
Arnoglossus tenuis	0	0	0	0	0	0	0	0	1	0	0	0	1
Cynoglossidae (tonguefish) Cynoglossus cf. brachycephalus	0	о	0	0	0	0	1	0	0	0	0	0	1
Cynoglossus macrolepidotus	0	0	0	0	0	0	Ĩ	0	0	0	0	0	1
Cynoglossus melampetalus	0	0	0	0	0	0	1	0	0	0	0	0	1
Cynoglossus puncticeps	1	0	0	0	0	0	0	2	0	0	0	0	3
Cynoglossus cf. puncticeps	. 0	0	0	0	0	0	0	0	0	1	0	0	1
Balistidae (filefishes & triggerfis Monacanthus nipponensis	hes) 0	0	0	0	1	0	0	1	1	0	0	0	3
Batrachoididae (toadfishes) <u>Batrachus grunniens</u>	0	0	0	11	1	0	0	0	0	1	0	0	13
Total specimens	55	5	4	29	14	56	12	23	17	12	3	10	240

.

VUNG-TAU (nondefoliated) -- dry season

VND stations	100	84	104	82	86	90	88	102	92	93	94	96	Total
Bottom salinity,	°/00 30,5	30.5	30.7	30.7	30.7	30.9	31,1	31.7	32,1	32. ;	33.3	33.5	
<u>Fishes</u> Clupeidae (herrings) ? <u>Sardinella</u> sp.	9	0	0	0	0	0	0	0	0	0	-	0	9
Engraulidae (anchovies) <u>Anchoviella</u> ? <u>tri</u>	0	0	0	0	0	0	0	0	0	2	-	o	2
Polynemidae (threadfins) <u>Polynemus</u> sp.	0	0	0	0	0	0	0	0	0	0	-	1	1
Centropomidae (snooks) <u>Ambassis</u> <u>dayi</u>	0	0	0	0	0	0	0	0	0	0	-	1	1
Apogonidae (cardinalfishes) Apogon quadrifasciatus	0	_ 0	0	1	0	0	0	5	0	0	-	0	6
Leiognathidae (ponyfishes) Leiognathus insidiator	7	4	0	0	0	1	0	0	1	1	-	0	14
Gerreidae (mojarras) Gerres filamentosus	2	1	1	Ø	0	0	0	1	0	0	-	0	5
Pomadasyidae (grunts) Pomadasys hasta	0	1	5	0	0	70	1	1	5	0	-	35	118
Scisenidae (croakers) Argyrosomus aneus	0	0	0	0	0	0	0	0	1	0	-	0	1
<u>Nibea</u> ? <u>albiflora</u>	0	0	0	0	0	0	0	0	0	0	-	9	9
<u>Sciaena russelli</u>	0	0	0	0	0	1	0	0	0	1	-	0	2
?Wak sina	1	0	0	0	0	0	2	0	0	1	-	56	60

٠

-

-

86

٠

VUNG-TAU (nondefoliated) -- dry season - (cont'd.)

VND stations	100	84	104	82	86	90	88	102	9 2	9 í	9 4	96	Total
Bottom salinity, ⁰ /00	30.5	30.5	30.7	30.7	30.7	30.9	31.1	31.7	32.1	32.3	33.3	33.5	
<u>Fishes</u> Callionymidae (dragonets) <u>Calliurichthys</u> ?fluviatilis	0	0	0	٥	0	0	0	2	0	0	-	0	2
Eleotridae (sleepers) Prionobutis koilomatodon	0	2	0	1	0	0	0	0	0	0	•	0	3
Gobiidae (gobies) ? <u>Acentrogobius</u> caninus	0	0	0	0	0	0	0	1	Ũ	0	-	0	1
?Cryptocentrus cf. russus	0	6	0	0	0	0	0	0	0	0	-	0	6
Unidentified genus	1	0	1	0	0	0	0	0	0	0	-	1	3
Scorpsenidae (scorpionfishes) ? <u>Prosopodasys</u> sp. ?Minous monodactylus	0	0	0	0	0	0	0	1	0	0	-	0	1
Platycephalidae (flatheads) Grammoplites scaber	0	0	. 0	0	0	0	0	2	1	0	-	1	2
Balistidae (filefishes & triggerfis <u>Monacanthus</u> <u>sulcatus</u>	shes) 0	0	0	0	0	0	0	1	0	0	-	0	1
Tetraodontidae (puffers) <u>Fugu</u> ?ocellatus	0	0	1	0	0	0	0	0	0	0	-	0	1
Total specimens	20	14	8	2	0	72	3	14	8	5	-	104	250

Table XIV.

Diversity indices of adult and juvenile fish families collected with bottom trawls in the estuaries of SVN. Stations are arranged in order of increasing salinity.

RUN	G-SAT SPECIAL ZO	DNE	v	UNG-TAU EGION	
VND stations	Wet season	Dry season	VND stations	Wet_season	Dry season
10	0.89	1.00	42	1.21	1.12
16	1,29	1.42	44	1.79	1,34
14	1,00	0.87	40	1.50	1.41
12	1,00	1.00	46	1.48	1.42
20	1,28	1.16	48	1.87	0.00
22	0.94	1.65	56	1.07	0,35
24	2,00	0.95	58	1.73	1.73
32	1.73	1.13	50	1.46	1.87
30	1.42	0.43	38	1.21	1.41
18	0,71	1.42	52	1,73	1.34
28	2,00	1.39	54	0.58	_a
34	0,00		36	1.90	0.59
26	1.50				
weighted average	1.67	0.68		1.10	1.01

⁸Net did not fish properly

.....

۰.

-

...

.

ţ,

Table XV.

1

Diversity indices of adult and juvenile fish species collected with bottom trawls in the estuaries of SVN. Stations are arranged in order of increasing bottom salinity.

<u>R</u>	UNG-SAT SPECIAL 2	CONE	<u>VI</u>	JNG-TAU REGION	
VND stations	Wet_season	Dry season	VND stations	Wet season	Dry season
10	0.89	1.00	42	0,13	1.12
16	1.81	1.42	44	1.79	1.34
14	1.00	0.87	. 40	1.50	1.41
12	1.00	1.00	46	1.67	1.42
20	1.49	1.45	48	2.14	0.00
22	1.42	1.42	56	0.13	0,35
24	2,50	1.95	58	2.31	1.16
32	1.73	1.13	4 50	1.67	2.14
30	1,42	0.51	38	1.21	1.41
18	0.71	1.42	52	1.73	1.79
28	2,00	1.66	54	0.58	- ^a
34	0.00		36	1.90	0.69
26	1.50				
weighted average	2.84	1.16		2.19	1.39

^a Net did not fish properly

greatest diversity of species. As with family diversity, species diversity was poorest in the Rung-SatSpecial Zone during the dry season. The richness of the Rung-Sat fish fauna during the wet season probably reflects the wide salinity regime encountered here.

These differences in part show the occurrence of a larger number of fish families, and thus may reflect a greater variety of fishes appearing in waters of higher salinity, rather than a lessened productivity of the defoliated region during the dry season. This increase in number and variety of fishes is presumably related to the greater variety of habitats found in salt water, and to the fact that the nonestuarine environment is less severe in its physiological and ecological demands.

In this particular case, however, we must consider the difficulty encountered in properly collecting juvenile and adult fishes. Because of the swift tidal currents usually encountered in the Rung-Sat Special Zone, it was sometimes not possible to determine if the net was always fishing properly on the bottom, and hence a smaller catch might have been obtained. Conversely, the comparatively cleater waters of the Vong Far Region might have permitted

fishes to detect visually the oncoming set of thes avoid it, a factor that also would reach in a smaller catch - Chelly, many kinds of fishes, especially the imper of swift-swimelow gets of one not easily captured with a small bettem trawl of the type models this study, necessitated by security and logistics. The only other type of sampling for large fishes entailed the use of MK3A2 hand grenades. Shallow, quiet-water embayments in both the Rung-Sat Special Zone and the Vung-Tau Region were frequently, though not statistically, sampled by grenades that

were set to explode at the bottom or just beneath the sea surface. Although juvenile and adult fishes were frequently obtained in the Vung-Tau Region using grenades, dead fishes were never observed at the surface in the Rung-Sat Special Zone following the use of grenades. Only fishes possessing air bladders will float, and, while it was considered <u>a priori</u> that possibly fishes having gas bladders were uncommon in the Rung-Sat Special Zone, the trawl net in this zone frequently captured juvenile fishes of families possessing gas bladders. We thus interpret this to indicate a real lack of juvenile and adult fishes in the areas sampled by grenade in the Rung-Sat Special Zone.

Only on two occasions in the Rung-Sat Special Zone did we observe fishes swimming. We could not identify the species in one instance, while in the other, at the southernmost limit of the Rung-Sat Special Zone, we observed a stingray leaping from the water. Conversely, in the Vung-Tau Region we frequently saw schools of mullet, anchovies, herring, silversides, and halfbeaks swimming or jumping at the surface. This indicates not only the greater clarity of the water in the Vung-Tau Region, which permitted us to see the fishes, but also the greater biological diversity of that area.

<u>Bird Take</u>. Although none of the personnel involved in this study was a trained ornithologist, whenever possible we attempted to note the presence and activity of birds in both study areas. Generally speaking, the Rung-Sat Special Zone had a greater number of insect-eating birds such as swallows. Sandpipers were occasionally observed feeding along the banks. A few ospreys (piscivores) and sea eaglets were observed here as well. In the Vung-Tau Region there were also kingfishers, a stork,

night herons, blue herons, terns, and gulls, in addition to the species observed in the Rung-Sat Special Zone. Birds were more frequently seen in the Vung-Tau Region and appeared to be more numerous. However, we were surprised at the relative lack of numbers and diversity of bird life along the beach and in the mangroves of the Vung-Tau Region in comparison with a somewhat similar yet larger and more speciose avifauna of South Florida.

The Food Web. The food and feeding relationships of organisms indicate interrelationships within an ecosystem (Odum 1971). Organisms are linked together by what they eat and what eats them. In a simple food chain, this relation tends to be linear, while in a more advanced ecosystem the interrelationships become complex webs of energy flow. Severe stress or rapid environmental changes involve reduction of the complex food web to simple forms involving only a few species (de Sylva 1973b).

To evaluate the food webs in the defoliated and nondefoliated areas, stomach contents of 323 juvenile and adult fishes were examined (Table XVI). These 46 species represent detritus feeders, planktivores, and primary, secondary, and tertiary carnivores feeding at all levels of the food web (Table XVII). Unfortunately, because our nets usually did not capture large fishes, knowledge of their food habits is limited. Nevertheless, juvenile and adult fishes of comparable size were caught in both study areas, and it appears reasonable to compare food webs of fishes of the same size in the two regions (Table XVIII).

Table XVI.

Number of stomachs containing food in fishes (2-28 cm) collected by 10-ft otter trawl in SVN. The Rung-Sat is defcliated; Vung-Tau is nondefoliated (control) region. Wet-season collections were made in October and November 1972. Dry-season collections were made in January 1973.

made in Decoper and No.	vember	. 191		Dry-	-seas	son cos	Lieca	lous	s wei	се ша	de 1	in Ja	111181	ry 19	973.						ped
	Food item	Nematoda	Polychaeta	Gastropoda	Bivalvia <u> </u>	Crustacea (uniden.)	Copepoda 60	Tsopoda	Amphipoda a	Decapoda a	Penaeidea	Caridea	Brachyura	Insecta Colectiera	Ophiuroides	Pisces	Algae or vege- table debria	Unidentified fleeb	Mud or sand	Empty stomachs	Number of fish stomachs exami
Fish species																					
Dasyatis ?kuhlii		0	0	0	0	0	1	0	1	1	0	0	0	0	0	0	0	0	0	0	1
Urolophoides sp.		0	0	0	0	0	0	0	0	2	1	2	1	0	0	1	0	1	0	1	3
<u>Coilia gravi</u>		0	0	0	0	0	6 [.]	0	0	1	0	0	0	0	0	0	0	0	0	0	6
Harpodon nehereus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1
Arius macronotacanthus		3	- 7	0	0	0	2	2	1	2	0	2	2	1	1	0	1	3	0	0	12
<u>Plotosus</u> canius		0	0	1	1	0	0	0	o	1	0	0	1	0	0	0	0	0	0	0	1
Polynemus longipectoralis		1	0	0	0	0	1	0	1	0	0	0	0	0	0	0	1	0	0	0	2
Lutjanus ?johni		0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	1
Leiognathus berbis		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Gerres abbreviatus		1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Pomadasys hasta		0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1
?Nibea acuta		1	0	0	1	0	3	1	2	3	0	0	2	0	0	0	1	1	0	1	5
Prionobutis koilomatodon		0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Acentrogobius caninus		1	0	0	0	0	0	0	0	0	0	0	0	0	·0	0	1	0	0	1	2

				10	1016	V A T	, 00	nem	ueu.												-
	RUNG	-5/	AT (d	lefo]	liate	ed)	W	et s	easo	0 -	(con	t'd.)				t 01 5	ø		8	sh amine
	Food item	Nematoda	Polychaeta	Gastropoda	Bivalvia	Crustacea	(uniden.) Copepoda	Isopoda	Amphipoda	Decapoda	(uniden.) Penaeidea	Caridea	Brachyura	Insecta	Conteopreta Cphiur des	Pisces	Algae or veg	Unidentified	Mud or sand	Empty stomac	Number of fi stomachs ex
Fish species																					
Apocryptodon cf. madurensis		1	0	0	0	0	-1	1	0	0	0	0	0	0	0	0	1	0	0	0	2
Ctenogobius sp.		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Glossogobius giurus		0	0	0	Û	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Trypauchen vagina		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Cynoglossus ?arel		2	2	0	0	0	0	0	2	0	0	0	2	0	0	0	0	0	0	0	2
Cynoglossus lingua		2	1	0	2	2	1	0	0	1	0	0	1	0	2	1	1	0	1	0	4
Zebrias sp.		1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	2
Batrachus grunniens		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2
Number of fish stomachs in which food items occurred	1	3	12	2	5	3	16	4	7	12	1	5	10	1	3	2	7	5	3	9	53

.

•.

RUNG-SAT (defoliated) -- dry season

						RU	NG-S	AT (defo	liat	ed)	d	ry s	easo	n										neđ
Food items		Foraminifera	Porifera	Hydrozoa	Nematoda	Polychaeta	Oligochaeta	Gastropoda	Bivelvia	Copepoda	Stomatopoda	ľsopoda	Amph 1 pod a	Decapoda	(unigen.) Caridea	Anomura	Brachyura	Pisces	Manmello	Algae or vege-	tapie georis Unidentified fleek	Egge	Mud or sand	Empty stomachs	Number of fish stomachs exami
<u>Fish</u> <u>species</u>											~~~		+					****							
<u>Coilia grayi</u>		0	0	0	1	0	0	0	0	22	8	0	б	19	0	0	7 ^æ	0	0	0	0	0	0	1	24
Arius macronotacanthus		0	0	0	1	11	1	1	10	1	1	4	6	4	1	0	3	0	ł	3	1	0	6	0	17
Polynemus longipectorali	8	0	0	0	0	0	0	0	0	0	I	0	0	1	0	0	0	0	0	0	0	0	0	0	2
Gerres filamentosus		0	0	0	0	1	0	0	0	1	0	0	0	0	0	° 0	0	0	0	0	0	0	0	0	1
Pseudosciaena crocea		0	0	0	0	0	0	0	0	2	0	0	2	2	0	0	0	0	0	0	0	0	0	0	2
<u>Scizena russelli</u>		0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
?Wak sina		1	1	2	18	16	0	0	2	14	0	1	69	36	0	I	2	2	0	20	8	ı	0	5	90
Parachaeturichthys polyn	епа	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Trypauchen vagina		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	2
Leptosynanceia asteroble	pa	0	0	0	0	0	٥	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Cynoglossus brachycephal	<u>us</u>	0	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	1
Cynoglossus lingus		0	0	0	0	0	0	0	1	1	0	0	2	0	0	0	0	0	0	1	0	0	0	1	4
Cynoglossus puncticeps		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Number of fish stomachs in which food items occurred		1	1	2	20	28	1	1	14	43	11	5	86	62	1	1	12	2	1	24	10	1	6	11	147

a Including 20ea

							VUNG	-TAU	(no	ondef	0110	ted)		wet	s ea	800)ed
	Food 1tem	Foraminifera	Nematoda	Polychaeta	Gastropoda	Bivalvia	Crustacea	(united)) Ostracoda	Copepoda	Cirrepedia	Stomatopoda	Lsopoda	Amphîpoda	Decapoda	Carldes	Brachyura	Ophiuroidea	Pisces	Algae or vege- ***1* debrie	Unider tified flesh	Wood fragments	Shell fragments	Gravel	Mud or sand	Empty stomachs	Number of fish stonachs examin
Fish species													*		+											
<u>Dasyatis</u> ? <u>kuhlii</u>		0	0	1	0	0	0	0	1	0	0	0	2	2	0	1	0	0	0	0	0	0	0	0	0	2
Ambassis dayi		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Apogon quadrifasciat	a	0	0	0	0	0	1	1	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	4
Sillago ?sihama		0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	0	0	1
Lutjanus ?johni		0	0	0	0	0	0	0	0	0	0	0	0	2	1	0	0	0	0	0	0	0	0	0	2	5
Pomadasys hasta		0	0	0	0	0	0	0	2	0	0	0	0	9	0	0	0	0	0	2	0	0	0	1	0	10
? <u>Wak</u> <u>sina</u>		0	3	1	0	0	0	0	3	0	0	0	0	2	0	0	2	0	0	0	0	ዑ	0	0	1	4
<u>Sciaena russelli</u>		0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	Q	1	0	0	1	0	0	0	1
Scatophagus argus		0	0	0	0	0	1	0	1	0	0	0	1	0	0	O	0	0	1	1	0	0	0	0	0	1
<u>Calliurichthys</u> ? <u>fluviatilis</u>		0	0	0	0	0	0	0	0	0	0	0	0	C	0	0	0	0	0	0 /	0	0	0	0	1	1
<u>Prionobutis</u> koilomatodon		0	2	2	0	C	1	2	3	2	0	1	3	5	2	0	2	1	6	0	0	0	0	1	2	16
Acentrogobius caninu	\$	0	3	1	1	0	0	0	1	0	0	0	0	2	0	0	0	0	2	0	0	0	0	0	1	4
Bethygobius fuscus		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Glossogobius giurus		0	0	1	0	0	0	0	3	0	0	0	5	12	0	0	0	1	0	0	1	0	0	1	7	23
Grammoplites scaber		0	0	0	0	0	0	0	1	- 0	0	0	0	8	0	0	0	0	0	2	0	0	1	1	0	8

•

•

Table XVI, continued.

.

.

.

VUNG-TAU (nondefoliated) -- wet season

96

-

.

٠

,

						V	UNG-1	TAU ((non	defo	liate	ed)		wet	seas	on -	(co	nt'ð	.)	L		æ	it 6			5	th Imined
	Rood i ftem		Foramintfera	Nematoda	Polychaeta	Gastropoda	Bivalvia	Crustaces	Ostracoda	Copepoda	Cirrepedia	Stomatopoda	Isopoda	Amphi poda	Decapoda	Caridea	Brachyura	Ophiuroidea	Pisces	Algae or vege rable datu 's	Unidentified fleeh	Wood fragment	Shell fragmen	Gravel	Mud or sand	Empty stomach	Number of fis stomachs exe
	<u>Fish</u> species																										
	Arnoglosus tenuis		0	0	0	0	0	0	0	0	0	0	0	0	0	2 ^t	0	0	0	0	0	0	0	0	0	0	1
	<u>Cynoglossus</u> cf. <u>brachycephalus</u>		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
	Cynoglossus macrolepidotus		0	0	Ĺ	0	1	0	0	0	0	0	0	0	0	0	0	0	0	o	0	0	0	0	0	0	1
	Cynoglossus puncticeps	: •	0	0	0	0	1	0	0	ì	0	0	0	0	1	0	Ð	0	0	1	0	0	0	0	0	0	2
71	Cynoglossus cf. puncticeps		0	0	0	0	0	0	0	0	0	0	0	0	0	Ó	0	0	0	0	1	0	0	0	1	0	1
	Batrachus grunniens		1	4	0	1	2	0	0	0	0	1	0	0	1	2	10	1	0	2	0	1	0	0	4	2	13
	Number of fish stomach in which food items occurred	15	1	13	8	2	4	3	3	18	2	1	1	12	45	7	11	5	2	14	6	2	1	1	9	20	101

^bIncluding larvae

`• •

VUNG-TAU (nondefoliated) -- dry season

	rood <u>item</u>	Colychaeta	Jastropoda	itvalvia	Crustacea (uniden.)	Copepoda	itomatopoda	kaphi poda	ecapoda (uniden.)	Jarídea	it achyur a)ph1uro1dea	lsces	Jnidentified flesh	Cmpty stomachs	Number of fish stomache exami
<u>Fish</u> species									#		-=	. <u>.</u> Y		5	==	
Anchoviella ?tri		0	1	0	0	1	0	0	2	0	2 ^a	0	0	0	0	2
Apogon guadrifasciata		0	0	0	0	1	0	0	2	0	0	0	1	0	3	6
Leiognathus insidiator		0	0	0	0	ò	0	0	0	0	0	0	0	1	1	2
Gerres filamentosus		2	0	0	0	3	0	3	1	0	0	0	0	0	0	4
Pomadasys hasta		0	0	0	0	1	0	2	1	0	0	0	0	0	I	3
Argyrosomus aneus		0	0	0	0	1	0	1	1	0	0	0	0	0	0	1
Nibea ?albiflora		0	0	0	2	4	0	3	0	0	0	1	0	2	0	5
Scigena russelli		0	0	0	0	2	0	0	0	0	0	0	0	0	0	2
?Wak sina		0	0	0	0	2	0	1	3	1	2 ⁸	0	0	0	1	6
<u>Calliurichthys</u> ? <u>fluviatilis</u>		0	0	0	0	0	0	0	0	0	0	0	0	0	1	1
Prionobutis koilomatodon		0	0	0	0	1	0	0	1	0	0	0	0	0	0	1
Grammoplites scaber		0	0	0	0	0	1	0	0	2	0	0	0	0	0	2
Monacanthus sulcatus		0	0	1	0	0	0	1	0	0	0	0	0	0	Ð	1
Fugu ? <u>ocellatus</u>		0	0	0	0	0	0	0	ο	0	0	0	0	0	1	1
Number of fish stomachs in which food items occurred		2	1	1	2	16	1	11	11	3	4	1	1	3	8	37

•

h mined

Table XVII.

Food habits of fishes in estuaries of SVN.

		Rung-Sat	(defolia	<u>Vung-Tau (nondefoliated)</u>								
	1972 wat season		19 dry s	73 eason	wet	1972 season	19 dry s	73 eason				
	No.	%	No.	%	No.	%	No.	%				
Foraminifera	0	0	1	1	1	· 1	0	0				
Sponges	0	0	1	1	0	0	0	0				
Hydrozoan s	0	0	2	1	0	0	0	0				
Nematodes	12	29	20	13	10	10	3	9				
Polychaetes	11	26	29	19	7	7	4	11				
Oligochaetes	1	2	0	0	0	0	0	0				
Gastropods	2	5	1	1	2	2	1	3				
Bivalves	3	7	13	9	3	3	1	3				
Crustacean uniden.	1	2	0	0	3	3	2	6				
Ostracods	2	5	0	0	3	3	0	0				
Copepods	16	38	43	29	15	16	16	46				
Cirrepedes	0	0	0	0	2	2	. 0	0				
Stomatopoda	0	0	12	8	1	1	1	3				
Isopoda	4	10	5	3	1	1	0	0				
Amphipods	8	19	86	58	14	15	8	23				
Decapods	11	26	65	44	42	44	12	34				

-

•

.

.

.

		Rung-Sat	t (defol:	lated)	Vung-Tau (nondefoliated)								
] wet	1972 season	dry	1973 season	wet	1972 season	dry	1973 season					
	No.	%	No.	%	No.	%	No.	%					
Peneids	1	2	0	0	0	0	0	0					
Carideans	5	12	1	1	6	6	3	9					
Anomura	0	0	1	1	0	0	0	0					
Brachyura	7	17	13	9	11	11	3	9					
Insects	1	2	0	0	0	0	0	0					
Ophiurians	1	2	0	0	3	3	3	9					
Fishes	1	2	2	1	2	2	1	3					
Mammals	Ó	0	1	1	0	0	0	0					
Algae	6	14	23	15	14	15	0	0					
Unidentified flesh	5	12	10	7	6	6	3	9					
Eggs	0	0	1	1	0	0	0	0					
Wood	٥	0	0	· 0	2	·2	0	0					
Shell	0	0	0	0	1	1	0	0					
Gravel	0	0	0	0	1	1	0	0					
Mud or sand	2	5	6	4	9	9	0	0					
Empty	5	12	10	7	19	20	8	23					
Total examined	42		150		96		35						

-

Table XVIII.

÷.

Lengths of fishes collected with 10-ft otter trawl (in mangrove region of SVN, 1972-73), which were examined for stomach contents. Lengths are fork lengths, except for Arnoglossus and Cynoglossus, for which total length was used. Size of Dasyatis and Urolophoides is disk width. ς, φ 11-11.9 ġ, 3-13.9 σ ch, 2-12. 22-22,5 9 φ, 6 δ 0-10. ľ. 23-23. œ۰, \$ \$ 6 4-14. 16-16. 15-15. -24 5-5. 8-8 6 5 e. 2-2. 6-6. 7-7. 4-4 Fish length (cm) ä j, Total -----Fishes examined 1 Desystis ?kuhlii 0 0 0 Ô 0 1 0 1 0 0 0 0 0 0 0 0 0 0 0 3 Urolophoides sp. 1 1 1 1 0 0 0 0 Ô Ð 0 0 1 0 0 5 Ö Ð Ô Ð O Anchoviella ?tri 2 n Ð n A a 0 Ð 2 0 ß a Ð £ A n 0 ก £ o 13 Coilia gravi 1 1 3 1 4 4 1 2 D 0 0 0 0 Û Û 0 Ô Ð 0 30 Harpodon nehereus Ð 0 0 1 0 0 Ð Ô ٥ 0 0 D Ð O 0 D n Û A 0 1 5 3 1 5 2 1 2 Ł Arius macronotacanthus 0 Ð 4 1 0 1 1 Ô O 1 n Ð 28 Plotosus canius 1 1 Q Ô. Ð Ð ٥ Ð 0 Ð 0 Ð 0 0 0 Ô 0 a 0 A 2 Polynemus longipectoralis 1 0 0 0 0 0 Û 1 0 0 0 0 0 0 0 2 0 0 0 0 0 Ambassis dayi 1 Ô n Ô. 0 0 0 0 0 0 0 0 0 0 a Ð Ð 0 0 1 Apogon quadrifasciata 3 1 1 4 1 Û 0 Û 0 10 0 0 0 O Û 0 0 0 0 a ٥ Sillago ?sihama 0 Ð 1 Ô 0 Ô 0 Ô 0 Ð Ó 0 Ô 0 0 0 0 0 O 0 1 0 1 Ð Û 0 0 0 0 0 Ð Lutjanus ?johni o 0 0 0 O 0 O 6 Leiognathus berbis 0 1 0 0 ٥ Û 0 n Ð 0 0 0 Ô Û 0 0 Ô Û 0 0 1 Leiognathus insidiator 0 0 0 0 0 0 0 0 0 1 0 1 0 0 0 0 0 0 0 0 2 Gerres abbreviatus 0 0 Ð 0 1 Û 0 Ð 0 0 0 0 Ô 0 0 0 0 0 0 1 0 Gerres filamentosus 0 S 2 0 o Ô Ð Ð Ð 0 a 0 ۵ Û, O £ A 0 0 7

101

1

· •

Fish length (cm)	2-2.9	3~3.9	4-4.9	5-5.9	6-6.9	7-7.9	8-8.9	6-9-9	10-10.9	11-11.9	12-12.9	13-13.9	14-14.9	15-15.9	16-16.9	17-17.9	22-22.9	23-23,9	24-24.9	27-27.9	Total
Fishes examined																					
Pomadasys hasta	2	7	2	0	0	0	0	1	0	0	0	0	0	0	0	Û	0	0	0	0	12
Argyrosomus abeus	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Nibes ?albiflora	0	2	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	5
?Nibea acuta	1	0	1	0	1	0	1	1	1	0	0	0	0	0	0	0	0	0	0	0	6
Pseudosciaena croces	0	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
<u>Sciaena russelli</u>	1	0	0	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	4
? <u>Wak sina</u>	2	16	17	14	18	14	10	6	O	2	0	1	0	1	0	0	0	0	0	0	101
Scatophagus argus	G	0	0	0	0	0	0	1	0	0	0	0	0	· 0	0	0	0	0	0	o	1
<u>Calliurichthys</u> fluviatilis	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Calliurichthys ?fluvistilis	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Prionobutis koilomatodon	2	7	5	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	18
Acentrogobius caninus	1	1	1	2	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	7
Apocryptodon cf. medurensis	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2
Bathygobius fuscus	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	ł
<u>Ctenogobius</u> sp.	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Glossogobius giurus	0	8	7	4	1	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	23
Parachaeturichthys polynema	0	0	1	0	Ō	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1

<u>Fish length (cm)</u>	2-2.9	3-3.9	4-4.9	5-5.9	6-6.9	7-7.9	8-8.9	9.9.9	10-10.9	11-11.9	12-12.9	13-13.9	14-14.9	15-15.9	16-16.9	17-17.9	22-22.9	23-23.9	24-24.9	27-27.9	Total
Fish examined	•				-																
Trypauchen vagina	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	. 3
Leptosynaceia asteroblepa	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Grammoplites scaber	0	Ð	0	0	3	1	0	0	0	3	1	0	0	0	0	2	1	0	0	0	11
Arnoglossus tenuis	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0.	0	0	1
Cynoglossus ?arel	0	0	0	0	0	1	2	0	0	0	0	0	0	0	0	0	0	0	0	0	3
Cynoglossus brachycephalus	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Cynoglossus cf. brachycephalus	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Cynoglossus lingua	0	1	0	0	0	2	2	0	2	0	0	0	0	0	0	0	1	0	1	0	9
Cynoglossus macrolepidotus	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1
Cynoglossus puncticeps	0	0	0	1	0	1	0	0	0	2	0	0	0	0	0	0	0	0	0	0	4
Cynoglossus cf. puncticeps	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
<u>Zebrias</u> sp.	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	Ð	0	0	0	0	2
Monacanthus sulcatus	0	0	1	0	0	0	0	0	Ð	0	0	0	0	0	0	0	0	0	0	0	1
Fugu ?ocellatus	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
Batrachus grunniens	0	0	0	5	1	5	0	1	0	0	0	1	1	0	1	0	0	0	0	0	15
Total specimens	10	63	47	42	45	37	27	15	8	22	6	4	6	3	3	2	2	1	1	1	345

103

.

---- , ,

During the wet season of the Vung-Tau Region (Figure 6), the food web is reasonably complex, showing several energy pathways and trophic levels that include secondary carnivores such as the flathead and the grunt. Detritus--an indication of a healthy, diverse ecosystem--is the prime source of energy. During the dry season (Figure 7), the pathways and relationships among the organisms are similar, but phytoplankton is believed to be an additional energy source, as evidenced by large quantities of planktonic copepods in surface waters.

In contrast the food web of the Rung-Sat Special Zone during the wet season is simple (Figure 8). In spite of the devastation of mangroves and the expected large amounts of detritus accumulating at the bottom of the estuaries, the strong tidal currents remove much wood, stems, and leaves of dead mangroves from these waters before they can decompose to organic detritus, and thus this energy is largely unavailable as a food source. Undoubtedly, however, the detritus is carried farther downstream where it is used for food in the littoral zone. During both the wet and dry seasons, high water turbidity in the Rung-Sat Special Zone probably reduces photosynthesis. Although diatoms occurred in both areas, they were far more abundant in the Vung-Tau Region, where turbidity is less (Table IV). During the dry season of the Rung-Sat Special Zone (Figure 9), a more complex and diverse food web is apparent, similar to that of the Vung-Tau Region during the dry season, except that detritus is the major energy source in the Rung-Sat Special Zone during the dry . season. With less runoff and higher salinities during the dry season,






FIG. 7. Food web in nondefoliated estuary, Vung-Tau, SVN, dry season.



FIG. 8. Food web in defoliated estuary, Rung-Sat Special Zone, SVN, wet season.

ł





a greater variety of organisms can invade the defoliated estuarine region and serve as food for predators. Yet secondary carnivores are scarce during both seasons, and probably reflect the stress ecosystem of the Rung-Sat.

There is no background data on food webs in Indochina. Since the country has been more or less continually at war since 1941, certainly many factors have affected the entire ecosystem. A hypothetical food web is depicted (Figure 10) that postulates energy pathways of man-made effects or war-related activities. The waters of the Saigon River, especially between Saigon and Mha-Be, contain a melange of sewage and industrial wastes, insecticides, paints, solvents, a variety of war materiel, explosives, chemicals, and waste products from hospitals flooded with war casualties. Along the river banks, war-time activities include, in addition to a variety of herbicides, the effects of napalm upon soils and vegetation, laterized soils from defoliation or Rome plowing, and rusting military hardware. We can only speculate upon the total effect of this combined effluent into the Rung-Sat Special Zone farther downstream.

The Commercial Marine Fisheries of Vietnam

The Vietnamese rely heavily upon aquatic life for food, with 80 to 90 percent of their protein coming from fish. Marine fish, crustaceans, and mollusks comprised 77 percent of the total fishery catch of Vietnam in 1969 (Brouillard 1970). That fishing is an important part of the Vietnamese economy is well documented (Soulier 1963, Loftas 1970).

During our studies in Vietnam, we were unable to participate in any



FIG. 10. Hypothetical energy pathways in the Rung-Sat estuaries, SVN, 1972.

trips aboard fishing vessels of any type. Frequent visits were made to fish markets around Saigon and at Vung-Tau and Xa-Thang, near Vung-Tau, to determine which species of fishes and invertebrates were commercially important. Some species were purchased for subsequent identification.

No major fishing activity was seen in the Rung-Sat Special Zone, probably because security still posed a problem, but possibly because this area is not (or perhaps never has been) a rich fishing area. Small boats were occasionally seen using a 2-poled dip net (<u>luci dan</u>), which pushed ahead of the boat. Presumably these nets catch anchovies, but we were not able to examine the catch. In the Vung-Tau Region, on the contrary, many such dip-net boats were observed. Common in this region are stake nets, which consist of numerous poles driven into the bottom, from which nets are strung. Fishes are enmeshed as they move with the tides, and are subsequently removed by the fishermen at the end of each tidal cycle. Also common were boats fishing drift gill nets and bottom set lines. Farther offshore, larger vessels used set lines, drift nets, shrimp trawls, and long gill nets.

Little detailed information is available on the number of pounds of shellfish (shrimps, crabs, oysters, clams) and marine fishes caught by commercial fishermen in SVN. Brouillard (1970) presented data on the marine fish catch of SVN from 1963 to 1969. It shows a gradual increase of from just under 3 million metric tons in 1963 to about 5 million metric tons in 1969 (Table XIX). This merely may indicate that the number of persons and boats fishing has increased.

Catch (metric tons), number of trawlers, and catch per unit of effort in the South Chira Ses and the Gulf of Thailand, 1952-1970. Data from Brouillard (1970) and Shindo (1973).

	Vietnamese vessels, coastal South Viet Nam			Japanese trawlers, South China Sea			Taiwanese trawlers, South China Sea			Thai trawlers, Gulf of Thailand		
	catch, tong x	travlers	effort	catch, tons	no. travlers	effort	catch, tons	no. baby trawler#	effortb	catch, tons x 103	no. trawlers	effort
1952	-	_8	-	3,079	14	219	-	-	-	-	-	-
1953	-	•	-	11,730	30	391	-	-	-	-	-	-
1954	-	-	-	12,045	31	388	2,214	468	4.3	-	-	-
1955	-	· -	-	8,171	23	355	2,623	558	7.8	-	-	-
1956	-	-	-	8,389	30	279	3,657	685	7.8	-	-	-
1957	• -	-	-	10,283	29.	354	4,802	867	6.9	-	•	-
1958	-	-	-	16,461	74	222	4,837	862	7.6	-	-	-
1959	-	-	-	8,155	23	354	5,518	918	8.8	-	-	.=
1960	-	-	-	5,647	14	403	6,040	962	9.1	-	-	-
1961	-	-	-	1,007	10	101	7,122	1,044	9,1	123	201	612
1962	-	-	-	384	4	96	6,984	964	12.7	151	976	155
1963	299	9,220	32	166	2	83	8,529	1,195	14.2	277	2,026	136
1964	314	9,710	32	Ο,	ò	0	11,905	1,386	15.5	372	2,360	158
1965	289	12,240	24	0	0	0	13,666	1,501	20.8	393	2,396	282
1966	287	16,770	17	0.	O	0	16,857	1,756	28.2	449	2,695	166
1967	319	23,195	14	٥	0	0	23,310	1,979	32.0	583	3,077	189
1968	321	29, 968	11	0	0	0	30,061	2,374	29.1	784	3,182	247
1969	355	39,001	9	0	0	0	32,886	2,278	33.9	908	3,185	285
1970	-	-	•	O	0	0	34,751	2,129	40.8	-	3,114	-
										•		

^a data not available

b annual total per gross ton

ì

A more meaningful analysis can be made if the number of motorized fishing vessels is considered as units of fishing gear, and these are divided each year into the marine fish catch for that year, giving a rough index of catch per unit of fishing effort (Figure 11). This catch curve indicates that the catch per unit of effort for marine fish species has steadily decreased, and shows a classic picture of a fishery that is steadily disappearing.

Historically, Japanese trawlers working in the South China Sea experienced considerable fluctuations in catch per effort (Shindo 1973). Between 1960 and 1961, the catch and catch per effort dropped drastically, indicating that overfishing by these vessels was already occurring. According to Shindo,

After 1958, the number of Japanese trawlers operating in the South China Sea would have increased rapidly, but the escalation of the war in Viet Nam prevented this development. Moreover, the decline of the catch rate in these waters made them less attractive. Hence the fishing operations ceased in 1963.

Because Thai trawlers fishing in the Gulf of Thailand are probably fishing many of the same racial stocks as occur in the South China Sea, a remarkably parallel drop in catch per effort of Thai trawlers is not unexpected.

It seems likely that Japanese trawlers were at least partly responsible for the abrupt decline in catch after 1960. The rate of decrease in catch per effort of Vietnamese trawlers is quite similar. With the cessation of extensive Japanese trawling operations, it would be expected that the Vietnamese catch per effort would have increased or at least stabilized if no factors other than fishing were operative. However,



FIG. 11. Catch per unit effort of fishing trawlers in the South China Sea, coastal SVN, and the Gulf of Thailand. Quantity and acreage of SVN sprayed with herbicides in from 1962 to 1970.

the ecological or socioeconomic effects of the war may have contributed to this decrease in effort, in spite of an increase in total fin-fish catch (Table X). In contrast, Thai trawlers working in areas of the Gulf of Thailand and the Taiwanese trawlers fishing the South China Sea, both of which are not affected by wartime activities, show a steady annual increase in the catch and the catch per unit of effort.

Loftas (1970) reported that "the size of fish caught [by Vietnamese fishermen] has decreased until the more popular species have had to be protected by setting size limits."

The decrease in the marine fish catch may also result from general pollution from domestic sewage, war materiel, or extensive spraying of DDT and malathion in Indochina, either alone or in combination with mangrove defoliants. None of these pollutants has been investigated in the aquatic food chain in SVN, but undoubtedly the chain is complexly affected. The somewhat sudden drop in the catch per unit effort after 1964 suggests that it is due not to overfishing by fishermen, but rather to an external factor such as environmental degradation or to some war-related sociological reason.

We lack supporting data to indicate whether or not the decline in the fishery is due to economic or sociological reasons related to the war. Nor is it known if the data presented on the number of motorized vessels used for catching marine fish indicate that these vessels are actually fishing full time, part time, or are even fishing in the same area from one day to the next. Possibly, for example, fuel or adequately trained fishermen are not always available to the

motorized fishing fleet. Indeed, certain parts of the coast of SVN are closed completely to fishing for security reasons, while in other sectors fishing is permitted only during certain hours. In some instances, vessels must fish a specified number of kilometers from shore. The added time required to run farther from port would decrease the time actually spent in fishing. Data on these factors are needed to evaluate possible causes and effects of the apparent decline in the fishing catch.

Security regulations on fishing activities plus restrictions following the Tet offensive were not imposed until 1968, whereas the decline in the marine fish catch per unit effort began abruptly in 1964. Extensive spraying of herbicidal defoliants began in 1963, following their limited use in 1961 and 1962 (Tables II and III, Whiteside 1970). One would expect a lag effect between initial widespread use of defoliants and effects on the fishery. It is tempting to compare the decrease in catch per effort by Vietnamese trawlers with the increase in gallons of herbicide or in number of acres sprayed in SVN (Figure 11).

One of the most important fishing ports in SVN is Vung-Tau (Brouillard 1970). If it can be assumed that in the Vung-Tau fishing area the fishing catch per unit effort decreased proportionally with the overall decrease in the catch per effort for the total marine fish catch in SVN, it can be stated biologically that most species brought into Vung-Tau and observed by us in the fish markets there are dependent upon estuarine conditions for food, shelter, or as

a nursery area. These include anchovies, herrings, flounders, snappers, croakers, porgies, jacks, whitings, mullets, shrimps, crabs, and prawns. Possibly, fishes and invertebrates landed at other South Vietnamese ports, at least south of Cam-Ranh, and east of Rach-Gia, at the Ca-Mau Peninsula, are also dependent upon the Rung-Sat Zone. Hence degradation of the Kung-Sat Special Zone would be expected to result in a decrease in the number of fish that depend upon the estuary for food, shelter, or as a nursery ground.

CONCLUSIONS

On the basis of very short-term studies made in March 1972, October and November 1972, and in January 1973, it may be concluded that defoliation of the mangrove area of the Rung-Sat Special Zone probably did not have a permanently damaging effect on the estuarine ecology of SVN. Biological productivity appears to be sufficiently high, although far from optimum, to permit the survival and feeding of a variety of organisms living in the estuary. It is likely, however, that this zone may have been much more productive prior to defoliation of the mangroves. There is little extensive spawning activity of fishes in the Rung-Sat Special Zone, probably because of the low oxygen saturation of the silt-laden, turbid waters. Nevertheless, larval fishes carried in from downstream are apparently able to survive under these conditions. One concern is that extensive defoliation between 1962-1970 may have resulted in a large "slug" of organic debris from grasses, leaves, twigs, and seeds, which is decomposing rather suddenly, resulting in a large volume of nutrients being added to the environment. Normal decay

of plant material and the conversion through detritus to the nutrient cycle should be retarded or prevented because of sudden defoliation (Bormann et al. 1968). This volume of nutrients would be abruptly used by phytoplankton and benthic algae or plants, resulting in "blooms." Yet it could be greater than the zooplankton-benthos mass that uses this large, sudden food supply. Nektonic or benthonic predators might not be numerous enough to use this sudden availability of energy, and thus this food source would not be utilized in the Rung-Sat Special Zone, though it could be assimilated farther downstream toward the Vung-Tau Region. In addition, after this sudden burst of food became available, nutrients would virtually disappear; since energy from mangrove degradation no longer existed, the normal detrital cycle would slow or cease. Hence, it may be anticipated that, unless other sources of nutrients are available in the Rung-Sat estuaries, the fishery will continue to decline.

The Rung-Sat Special Zone probably always was a difficult environment because of the wide tidal range and high tidal current velocities; only the latter might have changed appreciably as defoliation caused increased runoff from land. The increased organic silt load of the Rung-Sat Special Zone that probably followed defoliation is seen as the factor most damaging to the ecosystem of the Rung-Sat. Increased soil erosion following mangrove defoliation has resulted in the river filling with silt. The outcome is that there are fewer habitats for benthic fishes and invertebrates, and a correspondingly smaller area available to the commercial fishery. This siltation has also apparently increased the

abundance of tube-building polychaete worms, and has necessitated extensive dredging activities in the Saigon, Nha-Be, and Long-Tau Rivers; these projects add to the turbidity and siltation problem already occurring, destroying more benthic habitats. As long as siltation remains unchecked, turbidity will remain high in the rivers.

Another adverse effect of defoliation is that the resulting high turbidity of the waters in the Rung-Sat Special Zone may interfere with the photochemical degradation of 2,4-D and 2,4,5-T (Agent Orange) which leaches from the mangrove forests and soils of the Rung-Sat Special Zone. As a result, these compounds may persist in water, sediments, or aquatic organisms.

It has been suggested that herbicides might interfere with the production of phytoplankton; the relatively low numbers of diatoms found in the Rung-Sat Special Zone may be due to persistent defoliants, low light penetration, or both.

Because it was not possible to investigate their role and fate in the food web, it cannot be ascertained whether or not herbicides, their contaminants, or breakdown products were detrimental to the organisms in the ecosystem of the Rung-Sat Special Zone either by killing them directly or by interfering with physiological processes such as spawning, growth, swimming, behavior, and migrations. It is possible that the lower productivity of fish eggs, copepods, diatom, benthos, and fishes in the Rung-Sat Special Zone in comparison to that f the Vung-Tau Region was a result of residual herbicides in the water, rediment, or organisms. Similarly the role, pathways, and effects of hlorinated hydrocarbons,

organophosphates, heavy metals, and other contaminants entering the Rung-Sat Special Zone from upstream is unknown.

Finally, and possibly most important, the probability exists that if any of the above-mentioned contaminants enters the waters of the Rung-Sat Special Zone it will be incorporated into the food web, concentrated as it passes from one trophic level in the food web to each succeeding one, and eventually be passed on, in higher concentrations, because of biological magnification, to persons eating seafoods that have been associated with the food web of the Rung-Sat Special Zone.

RECOMMENDATIONS

1. This study suffered primarily from a lack of comparative background data. It is urged that the GVN and its academic institutions initiate a long-term ecological analysis, including biological, physical, and chemical observations and monitoring of the defoliated areas and aquatic ecosystem in SVN to evaluate recovery rates following defoliation of the mangrove ecosystem. The University of Miami's School of Marine and Atmospheric Science would be in a position to assist in the formulation of such programs, including the training of students and exchange between Vietnamese and American faculty and students.

2. Because fisheries development offers a real economic hope for Vietnam's future (Loftas 1970, Anonymous 1971), in conjunction with ecological studies it is recommended that the GVN institute a carefully-planned system of collecting accurate statistical data which would include, but not be limited to, catches of different species of fish and shellfish at each fishing port, number of boats

fishing on each fishing ground, time spent in actual fishing operations, kind of gear used, and economic data such as number of persons fishing. per boat, costs involved in ship operations, and cost of nets. A logbook system, such as is used in most other countries, would be desirable.

3. A program of replanting the mangroves of the Rung-Sat Special Zone should be initiated by planting seedlings or transplanting small trees. Special attempts should be made to plant the heavily eroded river banks with mangroves or other suitable plants that might reduce bank erosion and the resulting high turbidity of the Rung-Sat waters. Woodcutting activities should be prohibited close to the estuaries in order to maintain a buffer zone of mangroves and thus reduce erosion of river banks.

4. A thorough quantitative analysis of the food web of the Rung-Sat Special Zone is recommended, specifically to identify defoliants such as 2,4-D, 2,4,5-T, picloram, and cacodylic acid and their breakdown products and contaminants (dioxin and pentachlorophenol). Because dioxins also are found in paints, lacquers, and varnishes, sources of these should be searched for upstream of the Rung-Sat Special Zone. Other contaminants such as DDT and related chlorinated hydrocarbons and long-lived organophosphates should be identified, if they exist, in the food web. It would be desirable to identify and trace the pathways of all known contaminants through the water, sediments, organisms, and lower organisms. Finally, they may reach man.

ACKNOWLEDGMENTS

We wish to acknowledge the support of the National Academy of Sciences/National Research Council, Contract No. BA-23-73-47, to the University of Miami. We wish to thank various members of the NAS Committee on the Effects of Herbicides in Vietnam for their advice and assistance in the course of our field work in SVN, specifically Dr. Anton Lang, Committee Chairman, Professor Fham-Hoang-Ho, Professor Le-Van-Thoi, Dr. Fred Tschirley, Dr. Howard T. Odum, Dr. Joseph M. Carrier, and Dr. Philip Ross. We are grateful to Dr. Tran-ngoc-Loi and Mr. and Mrs. Nguyen-Thuong-Dao at the Institut Oceanographique at Nha-Trang, for much advice and assistance, and to Mr. Tran-Van-Tri and Mrs. Nam-Hai of the Department of Fisheries, Saigon. To many Vietnamese Naval personnel who guided and protected us during our field studies we are deeply grateful. Major Gordon Goff and Captain Dean Neptune, Military Assistance Command, Vietnam (MACV), were helpful in many ways.

We wish to thank our colleagues and coworkers at the University of Miami for various support during the field work in Vietnam and subsequent laboratory studies in Miami, including Helen Apuzzo, Maryanna Bogle, W. Connor Davis, Stephen Grund, Roger T. Hanlon, Eric Hildebrand, Raymond F. Hixon, James K. Low, John Markham, Susan McLean, Shirley Pomponi, James F. Quinn, Lewis N. Scotton, Robert C. Trumbull, Robert C. Work, and Bernard Yokel. We are grateful to Dr. Tokiharu Abe, Mrs. Lillian Dempster, Dr. William N. Eschmeyer, Dr. Carl L. Hubbs, and Ms. Terry Loy for assistance in locating pertinent literature.

Mrs. Selma Baron, of the National Academy of Sciences, assisted in many ways through obtaining necessary visas, arranging for shipping of materials, and general advice.

REFERENCES

Anonymous. 1971. South Vietnam looks to its fisheries. <u>Maritimes</u> 15:3-4.

Berry, A.J. 1964. Faunal zonation in mangrove swamps. <u>Bull. Natl.</u> Mus. Singapore 32:90-98.

Bormann, F.N., G.E. Likens, D.W. Fisher, and R.S. Pierce. 1968. Nutrient loss accelerated by clear-cutting of a forest. <u>Science</u> 159:882-4.

Brouillard, K.D. 1970. Fishery development survey: South Vietnam. Bureau of Commerical Fisheries, Department of the Interior, AID Project PIO/T 730-317-3-9190276, Saigon. 41 pp.

Davis, J.H. 1940. The ecological and geological role of mangroves in Florida. <u>Carnegie Inst. Washington Publ.</u> 517:303-412.

de Sylva, D.P. 1970. Ecology and distribution of postlarval fishes of southern Biscayne Bay, Florida. Progress Report to Environmental Protection Agency, Div. Water Qual. Res., Water Qual. Off. 198 pp.

. 1972. A visit to the mangrove region of South Viet Nam, March, 1972, with recommendation for future study. Report to the National Academy of Sciences/National Research Council Committee on the Effects of Herbicides in Vietnam. Univ. Miami, School of Marine and Atmospheric Science. 21 pp. (mimeogr.).

. 1973a. A bibliography pertinent to the marine and aquatic ecology of Viet Nam and adjacent areas. Unpublished manuscript.

. 1973b. Nektonic food webs in estuaries. <u>In</u> Recent advances in estuarine research, J.D. Costlow, ed. In press.

Erlich, P.R. and A.H. Erlich. 1972. Population; resources, environment: issues in human ecology, 2nd ed. W.H. Freeman, San Francisco. 509 pp.

Fell, J.W. and I.M. Master. 1973. Fungi associated with the degradation of mangrove (<u>Rhizophora mangle L.</u>) leaves in South Florida. <u>In</u> Belle W. Baruch Library, Marine Sci. I, Est. Micro. Biol. Ecol., Stevenson and Colwell, eds. Univ. South Carolina Press, Columbia. In press. Fischer-Piette, E. 1931. Sur la penetration des diverses especes marines sessiles dans les estuaires et sa limitation par l'eau douce. Ann. Inst. Oceanogr. Monaco 10:217-43.

Freise, F.W. 1935. A importancia da conservação dos mangroves como viveiros de peixes. An. 1^o Congr. Nac. de Pesca, Rio de Janeiro. pp. 315-9.

Freyberg, B.V. 1930. Zerstörung and Sedimentation an der Mangrovekuste Bradiliens. Leopoldina 6:69-118.

Gerlach, S.A. 1958. Die Mangroveregion tropischer Küsten als Lebensraum. <u>Z. Morphol.</u> <u>Ökol. Tiere</u> 46:636-730.

Giglioli, M.E.C. and I. Thornton. 1965. The mangrove swamps of Keneba, lower Gambia river basin: 1. Descriptive notes on the climate, the mangrove swamps and the physical composition of their soils. J. <u>Appl. Ecol.</u> 2:81-103.

Gledhill, D. 1963. The ecology of the Aberdeen creek mangrove swamp. <u>J. Ecol.</u> 51:693-703.

Golley, F.B., H.T. Odum, and R.F. Wilson. 1962. The structure and metabolism of a Puerto Rican red mangrove forest in May. Ecology 43: 9-19.

Heald, E.J. 1971. The production of organic detritus in a south Florida estuary. Univ. of Miami, Sea Grant Program. Tech. Bull. No. 6, Jan. 1971. 118 pp.

Hempel, G. 1965. On the importance of larval survival for the population dynamics of marine food fish. <u>Calif. Coop. Oceanic Fish.</u> <u>Invest. Rep.</u> 10:13-23.

Hynes, H.B.N. 1971. The biology of polluted waters. Liverpool Univ. Press, Liverpool. 202 pp.

Kato, K. 1966. Geochemical studies on the mangrove region of Cananeia Brazil: I. Tidal variations of water properties, II. Physiochemical observations on the reduction states. <u>Bol. Inst. Oceanogr.</u> 14:13-20, 21-24.

Kiener, A. 1966. Contributions a l'etude ecologique et biologique des eaux saumatres malagaches: les poissons euryhalins et leur role dans le developpement des peches. <u>Vie Milieu</u> 16:1013-1149.

Kobayashi, J. 1959. Chemical investigation on river waters of southeastern Asiatic countries. Report I: The quality of waters of Thailand. <u>Ber. Ohara Inst. Landwirtsch. Biol.</u> 11:167-233. Kopenski, R.P. 1968. Preliminary environmental data report, Saigon, Nha Be, Long Tau Rivers, Republic of Vietnam, 1967. U.S. Naval Oceanogr. Off., Washington. Unpaged.

Lauff, G.K., ed. 1967. Estuaries. Am. Assoc. Adv. Sci. Publ. 83. 757 pp.

Loftas, T. 1970. Fishery boom for South Vietnam. New Sci. 46:280-3.

Macnae, W. 1968. A general account of the fauna and flora of mangrove swamps and forests in the Indo-West-Pacific region. <u>In</u> Advances in marine biology, F.S. Russell and M. Yonge, eds. Vol. 6:73-270. Academic Press, New York. 406 pp.

Mansueti, R.J. 1962. Effects of civilization on striped bass and other estuarine biota in Chesapeake Bay and tributaries. <u>Proc. Gulf</u> Caribb. Fish. Inst., 14th Ann. Sess.:110-36.

Meselson, M.S., A.H. Westing, J.D. Constable, and R.E. Cook. 1970. Preliminary report. Herbicide Assessment Commission, AAAS. 8 pp. (mimeogr.).

Odum, E.P. 1971. Fundamentals of ecology. W.B. Saunders Co., Philadelphia. 574 pp.

Odum, W.E. 1970. Pathways of energy flow in a south Florida estuary. Univ. of Miami, Sea Grant Program. Tech. Bull No. 7. 173 pp. (Also RSMAS, Univ. of Miami, Ph.D. dissertation, 1970. 162 pp.)

and E.J. Heald. 1972. Trophic analyses of an estuarine mangrove community. Bull. Mar. Sci. 22:671-738.

Orians, G.H. and E.W. Pfeiffer. 1970. Ecological effects of the war in Vietnam. Science 168:544-54.

Shapley, D. 1973. Herbicides: AAAS study finds dioxin in Vietnamese fish. Science 180:285-6.

Shindo, S. 1973. General review of the trawl fishery and the demersal fish stocks of the South China Sea. FAO Fish. Tech. Paper 120: 1-49.

Shirota, A. 1966. The plankton of South Viet Nam. Overseas Tech. Coop. Agency, Japan. 462 pp.

Soulier, A. 1963. Preliminary bibliography of fish and fisheries for Cambodia, Laos, and Vietnam. <u>Indo-Pac. Fish. Counc. Occas. Pap.</u> 63(11). 80 pp.

Stephens, W.M. 1963. Mangroves: trees that make land. <u>Smithsonian</u> Inst. Annu. Rep. 1962;491-6.

Tabb, D.C., D.L. Dubrow, and R.B. Manning. 1962. The ecology of northern Florida Bay and adjacent estuaries. <u>Fla. State Board Conserv.</u> <u>Tech. Ser. 34:1-79.</u>

Tschirley, F.H. 1969. Defoliation in Vietnam. Science 163:779-86.

Walsh, G.E. 1967. An ecological study of a Hawaiian mangrove swamp. <u>In Estuaries</u>, G.H. Lauff, ed. AAAS, Washington. pp. 420-31.

Whiteside, T. 1970. Defoliation. Ballantine, New York. 168 pp.

N. W 1.